



Flg Monoclonal Antibody

Catalog No	YP-Ab-12925
Isotype	IgG
Reactivity	Human
Applications	WB;ELISA
Gene Name	FGFR1
Immunogen	Purified recombinant extracellular fragment of human Flg (aa22-376) fused with hlgGfc tag expressed in HEK293 cells.
Specificity	Flg Monoclonal Antibody detects endogenous levels of Flg protein.
Formulation	Ascitic fluid containing 0.03% sodium azide, 0.5% BSA, 50% glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	FGFR1; Basic fibroblast growth factor receptor 1; BFGFR; CEK; FGFR; FLG; FLT2; HBGR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus.
Tissue Specificity	Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.
Function	
Background	

**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images