



CRIF1 Polyclonal Antibody

Catalog No	YP-Ab-00360
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IHC;IF;ELISA
Gene Name	GADD45GIP1
Immunogen	The antiserum was produced against synthesized peptide derived from human GADD45GIP1. AA range:91-140
Specificity	CRIF1 Polyclonal Antibody detects endogenous levels of CRIF1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Growth arrest and DNA damage-inducible proteins-interacting protein 1;CRIF1; CR6-interacting;GADD45GIP1;PLINP1;PRG6;CKII beta-associating protein; CR6-interacting factor 1;Papillomavirus L2-interacting nuclear protein 1;PLINP; PLINP-1;p53-responsive
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Mitochondrion . Nucleus . Using N-terminally tagged constructs, has been found in the nucleus (PubMed:12482659). C-terminally tagged constructs are targeted exclusively to mitochondria (PubMed:22453275). This discrepancy may be explained by masking of a potential N-terminal mitochondrial targeting signal by the tag (PubMed:22453275). .
Tissue Specificity	Widely expressed. Highly expressed in the thyroid gland, heart, lymph nodes, trachea and adrenal tissues. Expressed at lower level in liver skeletal muscle, kidney, pancreas, testis, ovary and stomach. Barely detectable in adrenal adenoma and papillary thyroid cancer.
Function	function:Acts as a negative regulator of G1 to S cell cycle phase progression by inhibiting cyclin-dependent kinases. Inhibitory effects are additive with GADD45 proteins but occurs also in the absence of GADD45 proteins. Acts as a



repressor of the orphan nuclear receptor NR4A1 by inhibiting AB domain-mediated transcriptional activity. May be involved in the hormone-mediated regulation of NR4A1 transcriptional activity.,induction:Down-regulated by p53/TP53 in apoptotic cells.,miscellaneous:Cells overexpressing GADD45GIP1 were more likely to be in G1 and less likely to be in S phase and grow more slowly than control cells. Inhibiting the expression of GADD45GIP1 promotes cell cycle progression.,subunit:Interacts with GADD45A, GADD45B and GADD45G. Interacts with NR4A1 via the NR4A1 AB domain. Interacts with the human papilloma virus type 16 (HPV 16) minor capsid protein L2.,tissue specific

Background

This gene encodes a nuclear-localized protein that may be induced by p53 and regulates the cell cycle by inhibiting G1 to S phase progression. The encoded protein may interact with other cell cycle regulators. [provided by RefSeq, Aug 2012],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human brain, using GADD45GIP1 Antibody. The picture on the right is blocked with the synthesized peptide.