



Elongin A3 Polyclonal Antibody

Catalog No	YP-Ab-00387
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	TCEB3C
Immunogen	The antiserum was produced against synthesized peptide derived from human ELOA3. AA range:381-430
Specificity	Elongin A3 Polyclonal Antibody detects endogenous levels of Elongin A3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Elongin A3;TCEB3C;RNA polymerase II transcription factor SIII subunit A3; TCEB3L2;Elongin-A3;EloA3;Transcription elongation factor B polypeptide 3C
Observed Band	60kD
Calculated Molecular Weight	60kD
Cell Pathway	integral component of membrane,elongin complex,
Tissue Specificity	
Function	domain:The elongin BC complex binding domain is also known as BC-box with the consensus [APST]-L-x(3)-C-x(3)-[AIV].,function:SIII, also known as elongin, is a general transcription elongation factor that increases the RNA polymerase II transcription elongation past template-encoded arresting sites. Subunit A3 is transcriptionally active but its transcription activity is not enhanced by binding to the dimeric complex of the SIII regulatory subunits B and C (elongin BC complex) .,similarity:Contains 1 TFIIIS N-terminal domain.,subunit:Heterotrimer of an A (A1, A2 or A3), B and C subunit.,tissue specificity:Widely expressed.,

**Background**

The SIII (or elongin) transcription elongation factor complex stimulates the rate of transcription elongation by RNA polymerase II by suppressing the transient pausing of the polymerase at many sites along the DNA template. This complex is a heterotrimer, composed of the transcriptionally active subunit A, A2 or A3 (or elongin A, A2 or A3) and two regulatory subunits, B and C (or elongin B and C). This gene encodes subunit A3. A3 and A are ubiquitously expressed, whereas A2 is specifically expressed in the testis. [provided by RefSeq, Mar 2010],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images




Western Blot analysis of COLO205 cells using Elongin A3 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Western blot analysis of lysates from COLO205 cells, using ELOA3 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using CIDEA antibody.



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).