



ST5 Polyclonal Antibody

Catalog No	YP-Ab-00524
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	ST5
Immunogen	The antiserum was produced against synthesized peptide derived from human ST5. AA range:411-460
Specificity	ST5 Polyclonal Antibody detects endogenous levels of ST5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DENN domain-containing protein 2B;DENND2B;HTS1;ST5;Suppression of tumorigenicity 5 protein;HeLa tumor suppression 1
Observed Band	126kD
Calculated Molecular Weight	126kD
Cell Pathway	[Isoform 1]: Cytoplasm, cell cortex . Cell membrane . Recycling endosome . Colocalizes with RAB13 and ITSN1 at cytoplasmic vesicles that are most likely recycling endosomes. Colocalizes with the cortical actin cytoskeleton. .
Tissue Specificity	Widely expressed with the exception of peripheral blood lymphocytes. Isoform 1 is expressed in several epithelial and fibroblast (including tumorigenic) but absent in lymphoid cell lines (at protein level). Isoform 3 is expressed in primary cell or weakly tumorigenic but not in tumorigenic cell lines (at protein level).
Function	function:May be involved in cytoskeletal organization and tumorigenicity. Isoform 1 seems to be involved in a signaling transduction pathway leading to activation of MAPK1/ERK2. Isoform 3 may block ERK2 activation stimulated by ABL1. Isoform 3 may alter cell morphology and cell growth.,similarity:Contains 1 dDENN domain.,similarity:Contains 1 DENN domain.,similarity:Contains 1 uDENN domain.,subunit:Isoform 1 interacts with the SH3 domain of ABL1.,



tissue specificity:Widely expressed with the exception of peripheral blood lymphocytes. Isoform 1 is expressed in several epithelial and fibroblast (including tumorigenic) but absent in lymphoid cell lines (at protein level). Isoform 3 is expressed in primary cell or weakly tumorigenic but not in tumorigenic cell lines (at protein level).,

Background

This gene was identified by its ability to suppress the tumorigenicity of Hela cells in nude mice. The protein encoded by this gene contains a C-terminal region that shares similarity with the Rab 3 family of small GTP binding proteins. This protein preferentially binds to the SH3 domain of c-Abl kinase, and acts as a regulator of MAPK1/ERK2 kinase, which may contribute to its ability to reduce the tumorigenic phenotype in cells. Three alternatively spliced transcript variants of this gene encoding distinct isoforms are identified. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images

(kD)

Western Blot analysis of various cells using ST5
Polyclonal Antibody




Western Blot analysis of COLO205 cells using ST5
Polyclonal Antibody





Western blot analysis of lysates from COLO205 cells, using ST5 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using ST5 antibody.

