



SNAP 23 Polyclonal Antibody

Catalog No	YP-Ab-00764
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	SNAP-23
Immunogen	The antiserum was produced against synthesized peptide derived from the C-terminal region of human SNAP23. AA range:151-200
Specificity	SNAP 23 Polyclonal Antibody detects endogenous levels of SNAP 23 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HsT17016;SNAP 23;SNAP-23;SNAP23A;SNAP23B;Synaptosomal-associated protein 23;SNAP23;Vesicle-membrane fusion protein SNAP-23
Observed Band	25kD
Calculated Molecular Weight	25kD
Cell Pathway	Cell membrane; Peripheral membrane protein. Cell membrane; Lipid-anchor. Cell junction, synapse, synaptosome. Mainly localized to the plasma membrane.
Tissue Specificity	Ubiquitous. Highest levels where found in placenta.
Function	function:Essential component of the high affinity receptor for the general membrane fusion machinery and an important regulator of transport vesicle docking and fusion.,similarity:Belongs to the SNAP-25 family.,similarity:Contains 2 t-SNARE coiled-coil homology domains.,subcellular location:Mainly localized to the plasma membrane.,subunit:Binds simultaneously to SNAP25BP and SYN4. Found in a complex with VAMP8 and STX4 in pancreas (By similarity). Binds tightly to multiple syntaxins and synaptobrevins/VAMPs. Found in a complex with VAMP8 and STX1A.,tissue specificity:Ubiquitous. Highest levels where found in placenta.,

**Background**

Specificity of vesicular transport is regulated, in part, by the interaction of a vesicle-associated membrane protein termed synaptobrevin/VAMP with a target compartment membrane protein termed syntaxin. These proteins, together with SNAP25 (synaptosome-associated protein of 25 kDa), form a complex which serves as a binding site for the general membrane fusion machinery. Synaptobrevin/VAMP and syntaxin are believed to be involved in vesicular transport in most, if not all cells, while SNAP25 is present almost exclusively in the brain, suggesting that a ubiquitously expressed homolog of SNAP25 exists to facilitate transport vesicle/target membrane fusion in other tissues. The protein encoded by this gene is structurally and functionally similar to SNAP25 and binds tightly to multiple syntaxins and synaptobrevins/VAMPs. It is an essential component of the high affinity receptor for the

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images




Western Blot analysis of HepG2 cells using SNAP 23 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-ovary-cancer, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-ovary-cancer, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100



Western blot analysis of lysate from HepG2 cells, using SNAP23 Antibody.