



SERCA2 Polyclonal Antibody

Catalog No	YP-Ab-00772
Isotype	IgG
Reactivity	Human;Mouse;Rat;Chicken
Applications	WB;IHC;IF;ELISA
Gene Name	ATP2A2
Immunogen	The antiserum was produced against synthesized peptide derived from the C-terminal region of human ATP2A2. AA range:841-890
Specificity	SERCA2 Polyclonal Antibody detects endogenous levels of SERCA2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ATP2A2;SERCA2;SERCA2b;ATP2A 2;Sarcoplasmic/endoplasmic reticulum calcium ATPase 2;ATP2B;SR Ca(2+)-ATPase 2;Calcium pump 2;Calcium-transporting ATPase sarcoplasmic reticulum type;slow twitch skeletal muscle isoform;Endoplasmic reticulum class 1/2 Ca(2+) ATPase
Observed Band	115kD
Calculated Molecular Weight	115kD
Cell Pathway	Endoplasmic reticulum membrane ; Multi-pass membrane protein . Sarcoplasmic reticulum membrane ; Multi-pass membrane protein .
Tissue Specificity	Isoform 1 is widely expressed in smooth muscle and nonmuscle tissues such as in adult skin epidermis, with highest expression in liver, pancreas and lung, and intermediate expression in brain, kidney and placenta. Also expressed at lower levels in heart and skeletal muscle. Isoforms 2 and 3 are highly expressed in the heart and slow twitch skeletal muscle. Expression of isoform 3 is predominantly restricted to cardiomyocytes and in close proximity to the sarcolemma. Both isoforms are mildly expressed in lung, kidney, liver, pancreas and placenta. Expression of isoform 3 is amplified during monocytic differentiation and also observed in the fetal heart.



Function

alternative products: SERCA2 transcripts differ only in their 3'-UTR region and are expressed in a tissue-specific manner, catalytic activity: $\text{ATP} + \text{H}_2\text{O} + \text{Ca}^{2+}(\text{Cis}) = \text{ADP} + \text{phosphate} + \text{Ca}^{2+}(\text{Trans})$, disease: Defects in ATP2A2 are a cause of acrokeratosis verruciformis (AKV) [MIM:101900]; also known as Hopf disease. AKV is a localized disorder of keratinization, which is inherited as an autosomal dominant trait. Its onset is early in life with multiple flat-topped, flesh-colored papules on the hands and feet, punctate keratoses on the palms and soles, with varying degrees of nail involvement. The histopathology shows a distinctive pattern of epidermal features with hyperkeratosis, hypergranulosis, and acanthosis together with papillomatosis. These changes are frequently associated with circumscribed elevations of the epidermis that are said to resemble church spires. There are no feature

Background

This gene encodes one of the SERCA Ca^{2+} -ATPases, which are intracellular pumps located in the sarcoplasmic or endoplasmic reticula of muscle cells. This enzyme catalyzes the hydrolysis of ATP coupled with the translocation of calcium from the cytosol into the sarcoplasmic reticulum lumen, and is involved in regulation of the contraction/relaxation cycle. Mutations in this gene cause Darier-White disease, also known as keratosis follicularis, an autosomal dominant skin disorder characterized by loss of adhesion between epidermal cells and abnormal keratinization. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Oct 2008],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Products Images




Yao, Haidong, et al. "Selenoprotein W redox-regulated Ca²⁺ channels correlate with selenium deficiency-induced muscles Ca²⁺ leak." *Oncotarget* 7.36 (2016): 57618.


Western Blot analysis of 293 cells using SERCA2 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-muscle, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human testis. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human testis. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 30min).