



HAND1 Monoclonal Antibody

Catalog No	YP-Ab-00990
Isotype	IgG
Reactivity	Human
Applications	WB;ELISA
Gene Name	HAND1
Immunogen	Purified recombinant fragment of HAND1 (aa90-190) expressed in E. Coli.
Specificity	HAND1 Monoclonal Antibody detects endogenous levels of HAND1 protein.
Formulation	Ascitic fluid containing 0.03% sodium azide, 0.5% BSA, 50% glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	bHLHa27; Class A basic helix-loop-helix protein 27; eHAND; Extraembryonic tissues; heart; autonomic nervous system and; HAND1; Heart- and neural crest derivatives-expressed protein 1; autonomic nervous system and neural crest derivatives-expressed protein 1; eH
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Nucleus, nucleoplasm . Nucleus, nucleolus . Interaction with MDFIC sequesters it into the nucleolus, preventing the transcription factor activity. Phosphorylation by PLK4 disrupts the interaction with MDFIC and releases it from the nucleolus, leading to transcription factor activity (By similarity). .
Tissue Specificity	Heart.
Function	function: Plays an essential role in early trophoblast differentiation and in cardiac morphogenesis. In the adult, could be required for ongoing expression of cardiac-specific genes. Binds the DNA sequence 5'-NRTCTG-3' (non-canonical E-box)., similarity: Contains 1 basic helix-loop-helix (bHLH) domain., subunit: Efficient DNA binding requires dimerization with another bHLH protein. Forms homodimers and heterodimers with TCF3 gene products E12 and E47, HAND2 and HEY1, HEY2 and HEYL (hairy-related transcription factors)., tissue



specificity:Heart.,

Background

The protein encoded by this gene belongs to the basic helix-loop-helix family of transcription factors. This gene product is one of two closely related family members, the HAND proteins, which are asymmetrically expressed in the developing ventricular chambers and play an essential role in cardiac morphogenesis. Working in a complementary fashion, they function in the formation of the right ventricle and aortic arch arteries, implicating them as mediators of congenital heart disease. In addition, it has been suggested that this transcription factor may be required for early trophoblast differentiation. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis using HAND1 Monoclonal Antibody against truncated Trx-HAND1 recombinant protein (1).