



# Ikaros Monoclonal Antibody

|                                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Catalog No</b>                  | YP-Ab-01043                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>Isotype</b>                     | IgG                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Reactivity</b>                  | Human;Mouse;Dog                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Applications</b>                | WB                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Gene Name</b>                   | IKZF1                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Immunogen</b>                   | Purified recombinant human Ikaros (C-terminus) protein fragments expressed in E.coli.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Specificity</b>                 | Ikaros Monoclonal Antibody detects endogenous levels of Ikaros protein.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Formulation</b>                 | Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.2% sodium azide, 50% glycerol.                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| <b>Source</b>                      | Monoclonal, Mouse                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Purification</b>                | Affinity purification                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Dilution</b>                    | Western Blot: 1/1000 - 1/2000. Not yet tested in other applications.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Concentration</b>               | 1 mg/ml                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Purity</b>                      | ≥90%                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Storage Stability</b>           | -20°C/1 year                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| <b>Synonyms</b>                    | IKZF1;Ikaros;DNA-binding protein Ikaros;IK1;LYF1;ZNFN1A1;Ikaros family zinc finger protein 1;Lymphoid transcription factor LyF-1                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Observed Band</b>               |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Calculated Molecular Weight</b> |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Cell Pathway</b>                | Nucleus . In resting lymphocytes, distributed diffusely throughout the nucleus. Localizes to pericentromeric heterochromatin in proliferating cells. This localization requires DNA binding which is regulated by phosphorylation / dephosphorylation events. .; [Isoform Ik2]: Nucleus. In resting lymphocytes, distributed diffusely throughout the nucleus. Localizes to pericentromeric heterochromatin in proliferating cells. This localization requires DNA binding which is regulated by phosphorylation / dephosphorylation events (By similarity). .; [Isoform Ik6]: Cytoplasm . |
| <b>Tissue Specificity</b>          | Abundantly expressed in thymus, spleen and peripheral blood Leukocytes and lymph nodes. Lower expression in bone marrow and small intestine.                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| <b>Function</b>                    | function: Binds and activates the enhancer (delta-A element) of the CD3-delta gene. Functions in the specification and the maturation of the T-lymphocyte. Also interacts with a critical control element in the TDT (terminal deoxynucleotidyltransferase) promoter as well as with the promoters for other                                                                                                                                                                                                                                                                               |



genes expressed during early stages of B- and T-cell development., similarity:Belongs to the Ikaros C2H2-type zinc-finger protein family., similarity:Contains 6 C2H2-type zinc fingers.,subunit:Interacts with IKZF4 AND IKZF5.,tissue specificity:Abundantly expressed in thymus, spleen and peripheral blood Leukocytes and lymph nodes. Lower expression in bone marrow and small intestine.,

## Background

This gene encodes a transcription factor that belongs to the family of zinc-finger DNA-binding proteins associated with chromatin remodeling. The expression of this protein is restricted to the fetal and adult hemo-lymphopoietic system, and it functions as a regulator of lymphocyte differentiation. Several alternatively spliced transcript variants encoding different isoforms have been described for this gene. Most isoforms share a common C-terminal domain, which contains two zinc finger motifs that are required for hetero- or homo-dimerization, and for interactions with other proteins. The isoforms, however, differ in the number of N-terminal zinc finger motifs that bind DNA and in nuclear localization signal presence, resulting in members with and without DNA-binding properties. Only a few isoforms contain the requisite three or more N-terminal zinc motifs that confer high affinity binding


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis using Ikaros Monoclonal Antibody against Ramos, Jurkat cell lysate.