



Smad5 Monoclonal Antibody

Catalog No	YP-Ab-01051
Isotype	IgG
Reactivity	Human;Mouse;Rat;Bovine;Chicken;Pig;sheep
Applications	WB;IF;FCM
Gene Name	SMAD5
Immunogen	Purified recombinant human Smad5 (C-terminus) protein fragments expressed in E.coli.
Specificity	Smad5 Monoclonal Antibody detects endogenous levels of Smad5 protein.
Formulation	Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.2% sodium azide, 50% glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/1000 - 1/2000. Immunofluorescence: 1/100 - 1/500. Flow cytometry: 1/100 - 1/200. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MADH5;MAD homolog 5;JV5-1;JV5 1;hSmad5;Smad5;Mothers against decapentaplegic homolog 5;Mothers against DPP homolog 5;SMAD family member 5;SMAD 5
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Cytoplasm. Nucleus. Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4.
Tissue Specificity	Ubiquitous.
Function	function:Transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase. SMAD5 is a receptor-regulated SMAD (R-SMAD).,PTM:Phosphorylated on serine by BMP (bone morphogenetic proteins) type 1 receptor kinase.,PTM:Ubiquitin-mediated proteolysis by SMAD-specific E3 ubiquitin ligase SMURF1.,similarity:Belongs to the dwarfin/SMAD family.,similarity:Contains 1 MH1 (MAD homolog 1) domain.,similarity:Contains 1 MH2 (MAD homolog 2) domain.,subcellular location:Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4.,subunit:May form trimers with the co-SMAD SMAD4. Interacts with PEBP2-alpha subunit and



SMURF1. Interacts with SUV39H1 and SUV39H2.,tissue specificity:Ubiquitous.,

Background

The protein encoded by this gene is involved in the transforming growth factor beta signaling pathway that results in an inhibition of the proliferation of hematopoietic progenitor cells. The encoded protein is activated by bone morphogenetic proteins type 1 receptor kinase, and may be involved in cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2014],

matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images

Western Blot analysis using Smad5 Monoclonal Antibody against mouse testis lysate.



Immunofluorescence analysis of HeLa cells using Smad5 Monoclonal Antibody.



Flow cytometric analysis of Jurkat cells stained with Smad5 Monoclonal Antibody (red), followed by FITC-conjugated goat anti-mouse IgG. Black line histogram represents the isotype control, normal mouse IgG.

