



KCNK10 (TREK-2) Polyclonal Antibody

Catalog No	YP-Ab-01203
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;IHC;IF
Gene Name	KCNK10
Immunogen	Synthetic Peptide of KCNK10 (TREK-2) AA range: 16-66
Specificity	KCNK10(TREK-2) protein(A237) detects endogenous levels of KCNK10(TREK-2)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000, IHC 1:100-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	2P domain potassium channel TREK2;K2p10.1;KCNK10;KCNKA_HUMAN; Outward rectifying potassium channel protein TREK 2;Outward rectifying potassium channel protein TREK-2;Potassium channel subfamily K member 10; Potassium channel TREK 2;TREK 2;TREK 2 K;TREK 2 K(+) channel subunit; TREK-2 K(+) channel subunit;TREK2;TWIK related K+ channel 2;Potassium channel subfamily K member 10 (Outward rectifying potassium channel protein TREK-2) (TREK-2 K(+) channel subunit);Potassium channel subfamily K member 10 (Outward rectifying potassium channel protein TREK-2;TREK-2 K(+) channel subunit)
Observed Band	59kD
Calculated Molecular Weight	59kD
Cell Pathway	Membrane ; Multi-pass membrane protein .
Tissue Specificity	Abundantly expressed in pancreas and kidney and to a lower level in brain, testis, colon, and small intestine. Isoform b is strongly expressed in kidney (primarily in the proximal tubule) and pancreas, whereas isoform c is abundantly expressed in brain.



Function

function: Outward rectifying potassium channel. Produces rapidly activating and non-inactivating outward rectifier K(+) currents. Activated by arachidonic acid and other naturally occurring unsaturated free fatty acids., similarity: Belongs to the two pore domain potassium channel (TC 1.A.1.8) family., tissue specificity: Abundantly expressed in pancreas and kidney and to a lower level in brain, testis, colon, and small intestine. Isoform b is strongly expressed in kidney (primarily in the proximal tubule) and pancreas, whereas isoform c is abundantly expressed in brain.,

Background

potassium two pore domain channel subfamily K member 10(KCNK10) Homo sapiens The protein encoded by this gene belongs to the family of potassium channel proteins containing two pore-forming P domains. This channel is an open rectifier which primarily passes outward current under physiological K+ concentrations, and is stimulated strongly by arachidonic acid and to a lesser degree by membrane stretching, intracellular acidification, and general anaesthetics. Several alternatively spliced transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq, Sep 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of 1) Mouse Brain Tissue, 2) Rat Brain Tissue, 3) Jurkat with KCNK10(TREK-2) Rabbit pAb diluted at 1:2,000.



Immunohistochemical analysis of paraffin-embedded Rat Brain Tissue using KCNK10 (TREK-2) Rabbit pAb diluted at 1:200.