



GATA-1 (phospho Ser310) Polyclonal Antibody

Catalog No	YP-Ab-01244
Isotype	IgG
Reactivity	Human;Mouse;Rat;Monkey
Applications	WB;ELISA
Gene Name	GATA1
Immunogen	The antiserum was produced against synthesized peptide derived from human GATA1 around the phosphorylation site of Ser310. AA range:277-326
Specificity	Phospho-GATA-1 (S310) Polyclonal Antibody detects endogenous levels of GATA-1 protein only when phosphorylated at S310.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Anemia;X-linked;without thrombocytopenia;included;ERYF 1;Eryf1;Erythroid transcription factor;Erythroid transcription factor 1;GATA 1;GATA binding factor 1;GATA binding protein 1 (globin transcription factor 1);GATA binding protein 1;GATA-1;GATA-binding factor 1;GATA1;GATA1_HUMAN;GF 1;GF-1;GF1;Globin transcription factor 1;NF E1;NF E1 DNA binding protein;NF-E1 DNA-binding protein;NFE 1;NFE1;Nuclear factor erythroid 1;Transcription factor GATA1;XLANP;XLTD;XLTT
Observed Band	43kD
Calculated Molecular Weight	43kD
Cell Pathway	Nucleus.
Tissue Specificity	Erythrocytes.
Function	disease:Defects in GATA1 are the cause of X-linked dyserythropoietic anemia and thrombocytopenia (XDAT) [MIM:300367]. XDAT is a disorder characterized by erythrocytes with abnormal size and shape, and paucity of platelets in peripheral blood. The bone marrow contains abundant and abnormally small megakaryocytes.,disease:Defects in GATA1 are the cause of X-linked



thrombocytopenia with beta-thalassemia (XLTT) [MIM:314050]; also called thrombocytopenia, platelet dysfunction, hemolysis, and imbalanced globin synthesis. The disease consists of an unusual form of thrombocytopenia with beta-thalassemia. Patients have splenomegaly and petechiae, moderate thrombocytopenia, prolonged bleeding time due to platelet dysfunction, reticulocytosis and unbalanced (hemo)globin chain synthesis resembling that of beta-thalassemia minor.,domain:The two fingers are functionally distinct and cooperate to achieve

Background

This gene encodes a protein which belongs to the GATA family of transcription factors. The protein plays an important role in erythroid development by regulating the switch of fetal hemoglobin to adult hemoglobin. Mutations in this gene have been associated with X-linked dyserythropoietic anemia and thrombocytopenia. [provided by RefSeq, Jul 2008],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of various cells using Phospho-GATA-1 (S310) Polyclonal Antibody diluted at 1:500 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).



Western blot analysis of lysates from COS7 cells treated with EPO, using GATA1 (Phospho-Ser310) Antibody. The lane on the left is blocked with the phospho peptide.