



NFκB-p105 (phospho Ser907) Polyclonal Antibody

Catalog No	YP-Ab-01265
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;IP;ELISA
Gene Name	NFKB1
Immunogen	The antiserum was produced against synthesized peptide derived from human NF-kappaB p105/p50 around the phosphorylation site of Ser907. AA range:874-923
Specificity	Phospho-NFκB-p105 (S907) Polyclonal Antibody detects endogenous levels of NFκB-p105 protein only when phosphorylated at S907.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. Immunoprecipitation: 2-5 ug/mg lysate. ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DKFZp686C01211;DNA binding factor KBF1;DNA binding factor KBF1 EBP1; DNA-binding factor KBF1;EBP 1;EBP-1;EBP1;KBF1;MGC54151;NF kappa B; NF kappaB;NF kappabeta;NF kB1;NFkappaB;NFKB 1;NFKB p105;NFKB p50; Nfkb1;NFKB1_HUMAN;Nuclear factor kappa B DNA binding subunit;Nuclear factor kappa-B;subunit 1;Nuclear factor NF kappa B p105 subunit;Nuclear factor NF kappa B p50 subunit;Nuclear factor NF-kappa-B p50 subunit;Nuclear factor of kappa light chain gene enhancer in B cells 1;Nuclear factor of kappa light polypeptide gene enhancer in B cells 1;Nuclear factor of kappa light polypeptide gene enhancer in B-cells 1;p105;p50;p84/NF-kappa-B1 p98;Transcription factor NFKB1;Nuclear factor NF-kappa-B p105 subunit
Observed Band	110kD
Calculated Molecular Weight	110kD
Cell Pathway	Nucleus. Cytoplasm. Nuclear, but also found in the cytoplasm in an inactive form complexed to an inhibitor (I-kappa-B).
Tissue Specificity	Muscle,Rectum tumor,Uterus,



Function

domain:Glycine-rich region (GRR) appears to be a critical element in the generation of p50.,domain:The C-terminus of p105 might be involved in cytoplasmic retention, inhibition of DNA-binding, and transcription activation.,function:NF-kappa-B is a pleiotropic transcription factor which is present in almost all cell types and is involved in many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFkB1/p105, NFkB1/p50, REL and NFkB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Diff

Background

nuclear factor kappa B subunit 1(NFkB1) Homo sapiens This gene encodes a 105 kD protein which can undergo cotranslational processing by the 26S proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFkB) protein complex. NFkB is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NFkB translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NFkB has been associated with a number of inflammatory diseases while persistent inhibition of NFkB leads to inappropriate immune cell development or delayed cell growth. Alternative splicing results in multiple transcript variants encoding different isoforms

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images




Western Blot analysis of various cells using
Phospho-NFκB-p105 (S907) Polyclonal Antibody
diluted at 1:2000



Immunohistochemistry analysis of paraffin-embedded
human breast carcinoma, using NF-kappaB p105/p50
(Phospho-Ser907) Antibody.



NFkB-p105/p50 --
(pSer907)



-- 117

-- 85

-- 48

-- 34

-- 26

-- 19

(kD)

Western blot analysis of lysates from HeLa cells treated with TNF-alpha, using NF-kappaB p105/p50 (Phospho-Ser907) Antibody. The lane on the right is blocked with the phospho peptide.