



RUNX1 (phospho Ser249) Polyclonal Antibody


Catalog No	YP-Ab-01288
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	RUNX1
Immunogen	The antiserum was produced against synthesized peptide derived from human AML1 around the phosphorylation site of Ser276. AA range:242-291
Specificity	Phospho-RUNX1 (S249) Polyclonal Antibody detects endogenous levels of RUNX1 protein only when phosphorylated at S249.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Acute myeloid leukemia 1;Acute myeloid leukemia 1 protein;alpha subunit core binding factor;AML 1;AML1;AML1 EVI 1;AML1 EVI 1 fusion protein;Aml1 oncogene;AMLCR 1;AMLCR1;CBF alpha 2;CBF-alpha-2;CBFA 2;CBFA2;Core binding factor alpha 2 subunit;Core binding factor runt domain alpha subunit 2; Core-binding factor subunit alpha-2;EVI 1;EVI1;HGNC;Oncogene AML 1; Oncogene AML-1;OTTHUMP00000108696;OTTHUMP00000108697; OTTHUMP00000108699;OTTHUMP00000108700;OTTHUMP00000108702; PEA2 alpha B;PEA2-alpha B;PEBP2 alpha B;PEBP2-alpha B;PEBP2A2; PEBP2aB;Polyomavirus enhancer binding protein 2 alpha B subunit; Polyomavirus enhancer-binding protein 2 alpha B subunit;Run1;Runt related transcription factor 1;Runt-related transcription factor 1;RUNX 1;Runx1; RUNX1_HUMAN;SL3 3 enhancer factor 1 alpha B subunit;SL3-3 enhancer factor 1 alpha B subunit;SL3/AKV core binding factor alpha B subunit;SL3/AKV core-binding factor alpha B subunit;PEBP2-alpha
Observed Band	49kD
Calculated Molecular Weight	49kD
Cell Pathway	Nucleus.




Tissue Specificity	Expressed in all tissues examined except brain and heart. Highest levels in thymus, bone marrow and peripheral blood.
Function	alternative products:Additional isoforms seem to exist,caution:The fusion of AML1 with EAP in T-MDS induces a change of reading frame in the latter resulting in 17 AA unrelated to those of EAP.,disease:A chromosomal aberration involving RUNX1/AML1 is a cause of chronic myelogenous leukemia (CML). Translocation t(3;21)(q26;q22) with EAP, MSD1 or EVI1.,disease:A chromosomal aberration involving RUNX1/AML1 is a cause of chronic myelomonocytic leukemia. Inversion inv(21)(q21;q22) with USP16.,disease:A chromosomal aberration involving RUNX1/AML1 is a cause of M2 type acute myeloid leukemia (AML-M2). Translocation t(8;21)(q22;q22) with RUNX1T1/MTG8/ETO.,disease:A chromosomal aberration involving RUNX1/AML1 is a cause of therapy-related myelodysplastic syndrome (T-MDS). Translocation t(3;21)(q26;q22) with EAP, MSD1 or EVI1.,disease:A chromosomal aberration involving RUNX1/AML1 is found in child
Background	Core binding factor (CBF) is a heterodimeric transcription factor that binds to the core element of many enhancers and promoters. The protein encoded by this gene represents the alpha subunit of CBF and is thought to be involved in the development of normal hematopoiesis. Chromosomal translocations involving this gene are well-documented and have been associated with several types of leukemia. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of Jurkat cells using Phospho-RUNX1 (S249) Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).



Western blot analysis of lysates from Jurkat cells, using AML1 (Phospho-Ser276) Antibody. The lane on the right is blocked with the phospho peptide.