



C/EBP α (phospho Thr226) Polyclonal Antibody

Catalog No	YP-Ab-01302
Isotype	IgG
Reactivity	Human;Mouse;Rat;Monkey
Applications	WB;ELISA
Gene Name	CEBPA
Immunogen	The antiserum was produced against synthesized peptide derived from human C/EBP-alpha around the phosphorylation site of Thr226. AA range:192-241
Specificity	Phospho-C/EBP α (T226) Polyclonal Antibody detects endogenous levels of C/EBP α protein only when phosphorylated at T226.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	Apoptotic cysteine protease;Apoptotic protease Mch 5;C/EBP alpha;C/ebpalpha;CAP4;Caspase 8 precursor;CBF-A;CCAAT Enhancer Binding Protein alpha;CCAAT/enhancer binding protein (C/EBP);alpha;CCAAT/enhancer-binding protein alpha;CEBP;CEBP A;CEBP alpha;Cebpa;CEBPA_HUMAN;FADD homologous ICE/CED 3 like protease;FADD like ICE;FLICE;ICE like apoptotic protease 5;ICE8;MACH;MCH5;MORT1 associated CED 3 homolog;C/EBP α
Observed Band	42,also have 30kd isform
Calculated Molecular Weight	42,also have 30kd isform
Cell Pathway	Nucleus .; [Isoform 4]: Nucleus, nucleolus .
Tissue Specificity	Liver,Pancreas,Umbilical cord,White Matter pool- 5 brain tissues- f
Function	function:C/EBP is a DNA-binding protein that recognizes two different motifs: the CCAAT homology common to many promoters and the enhanced core homology common to many enhancers.,similarity:Belongs to the bZIP family.,similarity:Belongs to the bZIP family. C/EBP subfamily.,similarity:Contains 1 bZIP domain.,subunit:Binds DNA as a dimer and can form stable heterodimers with C/EBP beta and gamma. Interacts with UBN1. Interacts with HBV protein X.,

**Background**

This intronless gene encodes a transcription factor that contains a basic leucine zipper (bZIP) domain and recognizes the CCAAT motif in the promoters of target genes. The encoded protein functions in homodimers and also heterodimers with CCAAT/enhancer-binding proteins beta and gamma. Activity of this protein can modulate the expression of genes involved in cell cycle regulation as well as in body weight homeostasis. Mutation of this gene is associated with acute myeloid leukemia. The use of alternative in-frame non-AUG (GUG) and AUG start codons results in protein isoforms with different lengths. Differential translation initiation is mediated by an out-of-frame, upstream open reading frame which is located between the GUG and the first AUG start codons. [provided by RefSeq, Dec 2013],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Enzyme-Linked Immunosorbent Assay
(Phospho-ELISA) for Immunogen Phosphopeptide
(Phospho-left) and Non-Phosphopeptide
(Phospho-right), using C/EBP-alpha (Phospho-Thr226)
Antibody



Western blot analysis of lysates from COS7 cells
treated with EGF 200ng/ml 30', using C/EBP-alpha
(Phospho-Thr226) Antibody. The lane on the right is
blocked with the phospho peptide.