



# Smad3 (phospho Ser204) Polyclonal Antibody

|                                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Catalog No</b>                  | YP-Ab-01306                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Isotype</b>                     | IgG                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Reactivity</b>                  | Human;Mouse;Rat                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| <b>Applications</b>                | WB;IHC;IF;ELISA                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| <b>Gene Name</b>                   | SMAD3                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Immunogen</b>                   | The antiserum was produced against synthesized peptide derived from human Smad3 around the phosphorylation site of Ser204. AA range:170-219                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Specificity</b>                 | Phospho-Smad3 (S204) Polyclonal Antibody detects endogenous levels of Smad3 protein only when phosphorylated at S204.                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Formulation</b>                 | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Source</b>                      | Polyclonal, Rabbit,IgG                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Purification</b>                | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Dilution</b>                    | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Concentration</b>               | 1 mg/ml                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Purity</b>                      | ≥90%                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Storage Stability</b>           | -20°C/1 year                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Synonyms</b>                    | DKFZP586N0721;DKFZp686J10186;hMAD 3;hMAD-3;hSMAD3;HSPC193;HST17436;JV15 2;JV15-2;JV152;LDS1C;LDS3;MAD (mothers against decapentaplegic Drosophila) homolog 3;MAD homolog 3;Mad homolog JV15 2;Mad protein homolog;MAD;mothers against decapentaplegic homolog 3;Mad3;MADH 3;MADH3;MGC60396;Mothers against DPP homolog 3;SMA and MAD related protein 3;SMAD 3;SMAD;SMAD family member 3;Smad3;SMAD3_HUMAN                                                                                                                                                                    |
| <b>Observed Band</b>               | 48kD                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Calculated Molecular Weight</b> | 48kD                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Cell Pathway</b>                | Cytoplasm . Nucleus . Cytoplasmic and nuclear in the absence of TGF-beta. On TGF-beta stimulation, migrates to the nucleus when complexed with SMAD4 (PubMed:15799969, PubMed:21145499). Through the action of the phosphatase PPM1A, released from the SMAD2/SMAD4 complex, and exported out of the nucleus by interaction with RANBP1 (PubMed:16751101, PubMed:19289081). Co-localizes with LEMD3 at the nucleus inner membrane (PubMed:15601644). MAPK-mediated phosphorylation appears to have no effect on nuclear import (PubMed:19218245). PDPK1 prevents its nuclear |




translocation in response to TGF-beta (PubMed:17327236). Localized mainly to the nucleus in the early stages of embryo development with expression becoming evident in the cytoplasm of the inner cell mass at the blastocyst stage (By similarity)


|                                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|----------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Tissue Specificity</b>        | Brain,Colon carcinoma,Esophagus tumor,Pancreas,Placenta,Spleen,Umbilical cord blood                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Function</b>                  | disease:Defects in SMAD3 may be a cause of colorectal cancer (CRC) [MIM:114500].,domain:The MH2 domain is sufficient to carry protein nuclear export.,function:Transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinase. SMAD3 is a receptor-regulated SMAD (R-SMAD).,PTM:Phosphorylated on serine by TGF-beta and activin type 1 receptor kinases.,similarity:Belongs to the dwarfin/SMAD family.,similarity:Contains 1 MH1 (MAD homology 1) domain.,similarity:Contains 1 MH2 (MAD homology 2) domain.,subcellular location:In the cytoplasm in the absence of ligand. Migration to the nucleus when complexed with Smad4.,subunit:Interacts with HGS. Interacts with NEDD4L in response to TGF-beta. Interacts with TTRAP (By similarity). Interacts with SARA (SMAD anchor for receptor activation); form trimers with another SMAD3 and the co-SMAD SMAD4. Interacts wit |
| <b>Background</b>                | The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein functions as a transcriptional modulator activated by transforming growth factor-beta and is thought to play a role in the regulation of carcinogenesis. [provided by RefSeq, Apr 2009],                                                                                                                                                                                                                                                                                                                                                                                                              |
| <b>matters needing attention</b> | Avoid repeated freezing and thawing!                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Usage suggestions</b>         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |




## Products Images




Immunofluorescence analysis of Hela cell. 1, Smad3 (phospho Ser204) Polyclonal Antibody (red) was diluted at 1:200 (4° overnight). LC3A mouse Monoclonal Antibody (5G10) (green) was diluted at 1:200 (4° overnight). 2, Goat Anti Rabbit Alexa Fluor 594 Catalog: RS3611 was diluted at 1:1000 (room temperature, 50min). Goat Anti Mouse Alexa Fluor 488 Catalog: RS3208 was diluted at 1:1000 (room temperature, 50min).




Western Blot analysis of various cells using Phospho-Smad3 (S204) Polyclonal Antibody diluted at 1:500



Western Blot analysis of MCF-7 cells using Phospho-Smad3 (S204) Polyclonal Antibody diluted at 1:500



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Smad3 (Phospho-Ser204) Antibody