



FoxO4 (phospho Thr451) Polyclonal Antibody

Catalog No	YP-Ab-01312
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	FOXO4
Immunogen	The antiserum was produced against synthesized peptide derived from human FOXO4 around the phosphorylation site of Thr451. AA range:417-466
Specificity	Phospho-FoxO4 (T451) Polyclonal Antibody detects endogenous levels of FoxO4 protein only when phosphorylated at T451.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AFX;AFX1;Afxh;ALL1-fused gene from X chromosome;Fork head domain transcription factor AFX1;Forkhead box O4;Forkhead box protein O4;FOXO 4;Foxo4;FOXO4_HUMAN;MGC117660;MGC120490;Mixed lineage leukemia; translocated to;7;MLLT7;Myeloid/lymphoid or mixed lineage leukemia (trithorax homolog;Drosophila);Myeloid/lymphoid or mixed lineage leukemia;RGD1561201
Observed Band	55kD
Calculated Molecular Weight	55kD
Cell Pathway	Cytoplasm. Nucleus. When phosphorylated, translocated from nucleus to cytoplasm. Dephosphorylation triggers nuclear translocation. Monoubiquitination increases nuclear localization. When deubiquitinated, translocated from nucleus to cytoplasm.
Tissue Specificity	Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Isoform zeta is most abundant in the liver, kidney, and pancreas.
Function	disease:A chromosomal aberration involving FOXO4 is found in acute leukemias. Translocation t(X;11)(q13;q23) with MLL/HRX. The result is a rogue activator protein.,function:Transcription factor involved in the regulation of the



insulin signaling pathway. Binds to insulin-response elements (IREs) and can activate transcription of IGFBP1. Down-regulates expression of HIF1A and suppresses hypoxia-induced transcriptional activation of HIF1A-modulated genes. Also involved in negative regulation of the cell cycle.,pharmaceutical:A constitutively active FOXO4 mutant where phosphorylation sites Thr-32, Ser-187 and Ser-262 have been mutated to alanine may have therapeutic potential in ERBB2/HER2-overexpressing cancers as it inhibits ERBB2-mediated cell survival, transformation and tumorigenicity.,PTM:Acetylation by CBP, which is induced by peroxidase stress, inhibits transcriptional activity. Dea

Background

This gene encodes a member of the O class of winged helix/forkhead transcription factor family. Proteins encoded by this class are regulated by factors involved in growth and differentiation indicating they play a role in these processes. A translocation involving this gene on chromosome X and the homolog of the Drosophila trithorax gene, encoding a DNA binding protein, located on chromosome 11 is associated with leukemia. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2010],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Enzyme-Linked Immunosorbent Assay
(Phospho-ELISA) for Immunogen Phosphopeptide
(Phospho-left) and Non-Phosphopeptide
(Phospho-right), using FOXO4 (Phospho-Thr451)
Antibody



Western blot analysis of lysates from HUVEC cells
treated with EGF 200ng/ml 5', using FOXO4
(Phospho-Thr451) Antibody. The lane on the right is
blocked with the phospho peptide.