



FoxL2 (phospho Ser263) Polyclonal Antibody

Catalog No	YP-Ab-01315
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	FOXL2
Immunogen	The antiserum was produced against synthesized peptide derived from human FOXL2 around the phosphorylation site of Ser263. AA range:229-278
Specificity	Phospho-FoxL2 (S263) Polyclonal Antibody detects endogenous levels of FoxL2 protein only when phosphorylated at S263.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Blepharophimosis;Blepharophimosis epicanthus inversus and ptosis 1; Blepharophimosis epicanthus inversus and ptosis;BPES 1;BPES;BPES1; Epicanthus inversus and ptosis 1;Forkhead box L2;Forkhead box protein L2; Forkhead transcription factor FOXL2;FOX L2;FOXL 2;FOXL2;FOXL2_HUMAN; PFRK;PINTO;PITUITARY FORKHEAD FACTOR;POF 3;POF3
Observed Band	40kD
Calculated Molecular Weight	40kD
Cell Pathway	Nucleus .
Tissue Specificity	In addition to its expression in the developing eyelid, it is transcribed very early in somatic cells of the developing gonad (before sex determination) and its expression persists in the follicular cells of the adult ovary.
Function	disease:Defects in FOXL2 are a cause of blepharophimosis, ptosis, and epicanthus inversus syndrome (BPES) [MIM:110100]; also known as blepharophimosis syndrome. It is an autosomal dominant disorder characterized by eyelid dysplasia, small palpebral fissures, drooping eyelids and a skin fold running inward and upward from the lower lid. In type I BPSE (BPES1) eyelid

abnormalities are associated with female infertility. Affected females show an ovarian deficit due to primary amenorrhea or to premature ovarian failure (POF). In type II BPSE (BPES2) affected individuals show only the eyelid defects. There is a mutational hotspot in the region coding for the poly-Ala domain, since 30% of all mutations in the ORF lead to poly-Ala expansions, resulting mainly in BPES type II.,disease:Defects in FOXL2 are a cause of premature ovarian failure 3 (POF3) [MIM:608996]. Premature ovarian failure (POF)

Background

This gene encodes a forkhead transcription factor. The protein contains a fork-head DNA-binding domain and may play a role in ovarian development and function. Expansion of a polyalanine repeat region and other mutations in this gene are a cause of blepharophimosis syndrome and premature ovarian failure 3. [provided by RefSeq, Jul 2016],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Enzyme-Linked Immunosorbent Assay
(Phospho-ELISA) for Immunogen Phosphopeptide
(Phospho-left) and Non-Phosphopeptide
(Phospho-right), using FOXL2 (Phospho-Ser263)
Antibody



Western blot analysis of lysates from K562 cells treated
with Na₃VO₄ 0.3mM 40', using FOXL2
(Phospho-Ser263) Antibody. The lane on the right is
blocked with the phospho peptide.