



CREB-1 (phospho Thr100) Polyclonal Antibody

Catalog No	YP-Ab-01319
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	CREB1
Immunogen	The antiserum was produced against synthesized peptide derived from human CREB around the phosphorylation site of Thr100. AA range:71-120
Specificity	Phospho-CREB-1 (T100) Polyclonal Antibody detects endogenous levels of CREB-1 protein only when phosphorylated at T100.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Active transcription factor CREB;cAMP response element binding protein 1; cAMP response element binding protein;cAMP responsive element binding protein 1;cAMP-responsive element-binding protein 1;CREB;CREB-1;CREB1; CREB1_HUMAN;Cyclic AMP-responsive element-binding protein 1;MGC9284; OTTHUMP00000163864;OTTHUMP00000163865;OTTHUMP00000206660; OTTHUMP00000206662;OTTHUMP00000206667;Transactivator protein
Observed Band	42kD
Calculated Molecular Weight	42kD
Cell Pathway	Nucleus .
Tissue Specificity	Eye,Placenta,Spleen,Testis,
Function	disease:A chromosomal aberration involving CREB1 is associated with angiomatoid fibrous histiocytoma (AFH) [MIM:612160]. Translocation t(2;22)(q33;q12) with CREB1 generates a EWSR1/CREB1 fusion gene that is most common genetic abnormality in this tumor type.,function:This protein binds the cAMP response element (CRE), a sequence present in many viral and cellular promoters. CREB stimulates transcription on binding to the CRE. Transcription



activation is enhanced by the TORC coactivators which act independently of Ser-133 phosphorylation. Implicated in synchronization of circadian rhythmicity., PTM: Stimulated by phosphorylation. Phosphorylation of both Ser-133 and Ser-142 in the SCN regulates the activity of CREB and participates in circadian rhythm generation. Phosphorylation of Ser-133 allows CREBBP binding (By similarity). Phosphorylated upon DNA damage, probably by ATM or ATR., similarit

Background

This gene encodes a transcription factor that is a member of the leucine zipper family of DNA binding proteins. This protein binds as a homodimer to the cAMP-responsive element, an octameric palindrome. The protein is phosphorylated by several protein kinases, and induces transcription of genes in response to hormonal stimulation of the cAMP pathway. Alternate splicing of this gene results in several transcript variants encoding different isoforms. [provided by RefSeq, Mar 2016],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Enzyme-Linked Immunosorbent Assay
(Phospho-ELISA) for Immunogen Phosphopeptide
(Phospho-left) and Non-Phosphopeptide
(Phospho-right), using CREB (Phospho-Thr100)
Antibody



Western blot analysis of lysates from HUVEC cells
treated with etoposide 25uM 24H, using CREB
(Phospho-Thr100) Antibody. The lane on the right is
blocked with the phospho peptide.