



NF-κB2 p100 (phospho-Ser866/870) rabbit pAb

Catalog No	YP-Ab-01459
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	NFKB2 LYT10
Immunogen	Synthesized phosho peptide around human NF-κB2 p100 (Ser866 and 870)
Specificity	This antibody detects endogenous levels of Human Mouse NF-κB2 p100 (phospho-Ser866 or 870)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CVID10;DNA binding factor KBF2;H2TF1;Lymphocyte translocation chromosome 10 protein;LYT 10;NF κB2;NFKB p52/p100 subunit;Nuclear factor Kappa B subunit 2;Nuclear factor of kappa light polypeptide gene enhancer in B cells 2 (p49/p100);Nuclear factor of kappa light polypeptide gene enhancer in B cells 2;Oncogene Lyt 10;p100;Transcription factor NFKB2;NF-κB p100/p52; NFKB2 LYT10;NF-κB2 p100 (Ser866/870);Nuclear factor NF-kappa-B p100 subunit (DNA-binding factor KBF2) (H2TF1) (Lymphocyte translocation chromosome 10 protein) (Nuclear factor of kappa light polypeptide gene enhancer in B-cells 2) (Oncogene Lyt-10) (Lyt10) [Cleaved into: Nuclear factor NF-kappa-B p52 subunit]
Observed Band	100kD
Calculated Molecular Weight	100kD
Cell Pathway	Nucleus. Cytoplasm. Nuclear, but also found in the cytoplasm in an inactive form complexed to an inhibitor (I-kappa-B).
Tissue Specificity	Leukemia,Lymph,Thymus,



Function

disease: A chromosomal aberration involving NFKB2 is found in a case of B-cell non Hodgkin lymphoma (B-NHL). Translocation t(10;14)(q24;q32) with IGHA1. The resulting oncogene is also called Lym-10C alpha variant. disease: A chromosomal aberration involving NFKB2 is found in a cutaneous T-cell leukemia (C-TCL) cell line. This rearrangement produces the p80HT gene which encodes for a truncated 80 kDa protein (p80HT). disease: In B-cell leukemia (B-CLL) cell line, LB40 and EB308, can be found after heterogeneous chromosomal aberrations, such as internal deletions. domain: The C-terminus of p100 might be involved in cytoplasmic retention, inhibition of DNA-binding by p52 homodimers, and/or transcription activation. domain: The glycine-rich region (GRR) appears to be a critical element in the generation of p52. function: NF-kappa-B is a pleiotropic transcription factor which is present in almost

Background

nuclear factor kappa B subunit 2(NFKB2) Homo sapiens This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protein encoded by this gene can function as both a transcriptional activator or repressor depending on its dimerization partner. The p100 full-length protein is co-translationally processed into a p52 active form. Chromosomal rearrangements and translocations of this locus have been observed in B cell lymphomas, some of which may result in the formation of fusion proteins. There is a pseudogene for this gene on chromosome 18. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of 1 MCF-7 treated with LPS, 2 MCF7, using primary antibody at 1:1000 dilution. Secondary antibody (catalog#: RS23920) was diluted at 1:10000