



Smad2 (Phospho Ser255) rabbit pAb

Catalog No	YP-Ab-01490
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	SMAD2 MADH2 MADR2
Immunogen	Synthesized peptide derived from human Smad2 (Phospho Ser255)
Specificity	This antibody detects endogenous levels of Human,Mouse,Rat Smad2 (Phospho Ser255)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Drosophila;homolog of;MADR2;hMAD-2;HsMAD2;JV18;JV18-1;JV181;MAD; MAD homolog 2;MAD Related Protein 2;Mad-related protein 2;MADH2; MGC22139;MGC34440;Mother against DPP homolog 2;Mothers against decapentaplegic homolog 2;Mothers against decapentaplegic;2;Mothers against DPP homolog 2;OTTHUMP00000163489;Sma and Mad related protein 2;Sma- and Mad-related protein 2 MAD;SMAD 2;SMAD family member 2;SMAD; SMAD2;SMAD2_HUMAN;SMAD2 MADH2 MADR2;Smad2 (Phospho Ser255); Mothers against decapentaplegic homolog 2 (MAD homolog 2;hSMAD2)
Observed Band	55kD
Calculated Molecular Weight	55kD
Cell Pathway	Cytoplasm . Nucleus . Cytoplasmic and nuclear in the absence of TGF-beta. On TGF-beta stimulation, migrates to the nucleus when complexed with SMAD4 (PubMed:9865696, PubMed:21145499). On dephosphorylation by phosphatase PPM1A, released from the SMAD2/SMAD4 complex, and exported out of the nucleus by interaction with RANBP1 (PubMed:16751101, PubMed:19289081). Localized mainly to the nucleus in the early stages of embryo development with expression becoming evident in the cytoplasm at the blastocyst and epiblast stages (By similarity). .



Tissue Specificity Expressed at high levels in skeletal muscle, endothelial cells, heart and placenta.


Function embryonic axis specification, in utero embryonic development, formation of primary germ layer, mesoderm formation, peptide secretion, generation of a signal involved in cell-cell signaling, regionalization, transcription, regulation of transcription, DNA-dependent, regulation of transcription from RNA polymerase II promoter, RNA processing, protein complex assembly, protein amino acid phosphorylation, phosphorus metabolic process, phosphate metabolic process, cell surface receptor linked signal transduction, enzyme linked receptor protein signaling pathway, transmembrane receptor protein serine/threonine kinase signaling pathway, transforming growth factor beta receptor signaling pathway, SMAD protein complex assembly, intracellular signaling cascade, cell-cell signaling, zygotic determination of dorsal/ventral axis, gastrulation, pattern specification process, mesoderm development, heart

Background

matters needing attention Avoid repeated freezing and thawing!

Usage suggestions This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human uterus. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).