



CNOT7 Polyclonal Antibody

Catalog No	YP-Ab-01617
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;IHC;IF;ELISA
Gene Name	CNOT7
Immunogen	The antiserum was produced against synthesized peptide derived from human CNOT7. AA range:10-59
Specificity	CNOT7 Polyclonal Antibody detects endogenous levels of CNOT7 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	BTG1 binding factor 1;CAF 1;CAF1;CCR4 associated factor 1;CNOT7;hCAF 1; CCR4-NOT transcription complex subunit 7;BTG1-binding factor 1;CCR4-associated factor 1;CAF-1
Observed Band	36kD
Calculated Molecular Weight	36kD
Cell Pathway	Nucleus. Cytoplasm, P-body . NANOS2 promotes its localization to P-body. .
Tissue Specificity	Embryo,Lymph,Placenta,Prostate,Testis,
Function	function:Ubiquitous transcription factor required for a diverse set of processes. It is a component of the CCR4 complex involved in the control of gene expression., similarity:Belongs to the CAF1 family.,subunit:Interacts with TOB1.,
Background	CCR4-NOT transcription complex subunit 7(CNOT7) Homo sapiens The protein encoded by this gene binds to an anti-proliferative protein, B-cell translocation protein 1, which negatively regulates cell proliferation. Binding of the two proteins, which is driven by phosphorylation of the anti-proliferative protein, causes signaling events in cell division that lead to changes in cell proliferation associated with cell-cell contact. The encoded protein downregulates the innate



immune response and therefore provides a therapeutic target for enhancing its antimicrobial activity against foreign agents. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 1 and X. [provided by RefSeq, Apr 2016],

matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images





Western Blot analysis of various cells using CNOT7 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).



Western blot analysis of lysates from HeLa and HUVEC cells, using CNOT7 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HUVECcells using CNOT7 antibody.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Tris-EDTA, pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200(4° overnight.3, Secondary antibody was diluted at 1:200(room temperature, 45min).