



HDAC8 Polyclonal Antibody

Catalog No	YP-Ab-01768
Isotype	IgG
Reactivity	Human;Mouse;Rat;Monkey
Applications	WB;IHC;IF;ELISA
Gene Name	HDAC8
Immunogen	The antiserum was produced against synthesized peptide derived from human HDAC8. AA range:5-54
Specificity	HDAC8 Polyclonal Antibody detects endogenous levels of HDAC8 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Protein deacetylase HDAC8;Protein deacetylase HDAC8;HDAC 8;EC:3.5.1.98; EC:3.5.1.;HDAC8;Histone deacetylase 8;HDACL1;CDA07;HD8
Observed Band	42kD
Calculated Molecular Weight	42kD
Cell Pathway	Nucleus . Chromosome . Cytoplasm . Excluded from the nucleoli (PubMed:10748112). Found in the cytoplasm of cells showing smooth muscle differentiation (PubMed:15772115, PubMed:16538051). .
Tissue Specificity	Weakly expressed in most tissues. Expressed at higher level in heart, brain, kidney and pancreas and also in liver, lung, placenta, prostate and kidney.
Function	catalytic activity:Hydrolysis of an N(6)-acetyl-lysine residue of a histone to yield a deacetylated histone.;caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.;function:Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes., miscellaneous:Its activity is inhibited by trichostatin A (TSA) and butyrate, two



well known histone deacetylase inhibitors.,similarity:Belongs to the histone deacetylase family. Type 1 subfamily.,subcellular location:Excluded from the nucleoli.,subunit:Interacts with PEPB2-MYH11, a f

Background

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class I of the histone deacetylase family. It catalyzes the deacetylation of lysine residues in the histone N-terminal tails and represses transcription in large multiprotein complexes with transcriptional co-repressors. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2009],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images




Western Blot analysis of various cells using HDAC8
Polyclonal Antibody diluted at 1:1000



Western Blot analysis of 293-UV cells using HDAC8
Polyclonal Antibody diluted at 1:1000



Immunohistochemistry analysis of paraffin-embedded
human lung carcinoma tissue, using HDAC8 Antibody.
The picture on the right is blocked with the synthesized
peptide.



Western blot analysis of lysates from NIH/3T3, HeLa, and Jurkat cells, , using HDAC8 Antibody. The lane on the right is blocked with the synthesized peptide.