



HIF-3α Polyclonal Antibody

Catalog No	YP-Ab-01777
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	HIF3A
Immunogen	The antiserum was produced against synthesized peptide derived from human HIF-3α. AA range:305-354
Specificity	HIF-3α Polyclonal Antibody detects endogenous levels of HIF-3α protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HIF3 alpha;HIF 3A4;HIF 3A2;HIF 3A;HIF 3 alpha;HIF-3α;HIF3A;Hypoxia-inducible factor 3-α;HIF-3-α;HIF3-α;Basic-helix-loop-helix-PAS protein MOP7;Class E basic helix-loop-helix protein 17;bHLHe17;HIF3-α-1;Inhibitory PAS domain protein;IPAS;Member of PAS protein 7;PAS domain-c
Observed Band	72kD
Calculated Molecular Weight	72kD
Cell Pathway	nucleus,nucleoplasm,mitochondrion,cytosol,nuclear speck,
Tissue Specificity	Cerebellum,Embryo,Kidney,Ovary,Pancreas,Whole embryo,
Function	function:Involved in adaptive response to hypoxia. Suppresses hypoxia-inducible expression of HIF1A and EPAS1. Binds to core DNA sequence 5'-TACGTG-3' within the hypoxia response element (HRE) of target gene promoters. The complex HIF3A-ARNT activates the transcription of reporter genes driven by HRE. Isoform 4 has a dominant-negative function of inactivating HIF1A-mediated transcription. Isoform 4 attenuates the binding of HIF1A to hypoxia-responsive elements (HRE), thus inhibiting HRE-driven transcription. Hypoxia induces down-regulation of isoform 4, leading to activation of HIF1A in



hypoxia. Conversely, upon restoring normoxia, the expression of isoform 4 increases and thereby secure an inhibition of HIF1A activity. Isoform 4 may be a negative regulator of hypoxia-inducible gene expression in the kidney and may be involved in renal tumorigenesis. Functions as an inhibitor of angiogen

Background

hypoxia inducible factor 3 alpha subunit(HIF3A) Homo sapiens The protein encoded by this gene is the alpha-3 subunit of one of several alpha/beta-subunit heterodimeric transcription factors that regulate many adaptive responses to low oxygen tension (hypoxia). The alpha-3 subunit lacks the transactivation domain found in factors containing either the alpha-1 or alpha-2 subunits. It is thought that factors containing the alpha-3 subunit are negative regulators of hypoxia-inducible gene expression. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Mar 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysate from rat brain cells, using HIF-3α antibody.