



Nopp140 Polyclonal Antibody

Catalog No	YP-Ab-01911
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	NOLC1
Immunogen	Synthesized peptide derived from Nopp140 . at AA range: 620-700
Specificity	Nopp140 Polyclonal Antibody detects endogenous levels of Nopp140 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HCV NS5A-transactivated protein 13;Hepatitis C virus NS5A-transactivated protein 13;KIAA0035;NOPP130;Nopp140;NOLC1;Nucleolar and coiled-body phosphoprotein 1;NS5ATP13;140 kDa nucleolar phosphoprotein;Nucleolar 130 kDa protein;Nucleolar pho
Observed Band	74,130KD(Nucleolar phosphoprotein p130)
Calculated Molecular Weight	74,130KD(Nucleolar phosphoprotein p130)
Cell Pathway	Nucleus, nucleolus . Cytoplasm . Shuttles between the nucleolus and the cytoplasm. At telophase it begins to assemble into granular-like pre-nucleolar bodies which are subsequently relocated to nucleoli at the early G1-phase. .
Tissue Specificity	Bone marrow,Epithelium,Leukemia,Liver,Lung,Placenta,T-cell,
Function	function:Related to nucleogenesis, may play a role in the maintenance of the fundamental structure of the fibrillar center and dense fibrillar component in the nucleolus. It has intrinsic GTPase and ATPase activities. May play an important role in transcription catalyzed by RNA polymerase I.,PTM:Undergoes rapid and massive phosphorylation/dephosphorylation cycles on CK2 and PKC sites. There is evidence suggesting that CDC2 kinase phosphorylates p130 at the M-phase.,similarity:Contains 1 LisH domain.,subcellular location:Shuttles between



the nucleolus and the cytoplasm. At telophase it begins to assemble into granular-like pre-nucleolar bodies which are subsequently relocated to nucleoli at the early G1-phase.,subunit:Interacts with RNA polymerase I 194 kDa subunit (RPA194) and with casein kinase-II.,

Background

function:Related to nucleogenesis, may play a role in the maintenance of the fundamental structure of the fibrillar center and dense fibrillar component in the nucleolus. It has intrinsic GTPase and ATPase activities. May play an important role in transcription catalyzed by RNA polymerase I.,PTM:Undergoes rapid and massive phosphorylation/dephosphorylation cycles on CK2 and PKC sites. There is evidence suggesting that CDC2 kinase phosphorylates p130 at the M-phase.,similarity:Contains 1 LisH domain.,subcellular location:Shuttles between the nucleolus and the cytoplasm. At telophase it begins to assemble into granular-like pre-nucleolar bodies which are subsequently relocated to nucleoli at the early G1-phase.,subunit:Interacts with RNA polymerase I 194 kDa subunit (RPA194) and with casein kinase-II.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of various lysis using Nopp140 Polyclonal Antibody diluted at 1:2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000