



TFIIH p44 Polyclonal Antibody


Catalog No	YP-Ab-02106
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	GTF2H2
Immunogen	The antiserum was produced against synthesized peptide derived from human TF2H2. AA range:1-50
Specificity	TFIIH p44 Polyclonal Antibody detects endogenous levels of TFIIH p44 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	BTF2;BTF2 p44;BTF2P44;T BTF2P44;TFIIH;TFIIH p44;GTF2H2;General transcription factor IIH subunit 2;Basic transcription factor 2 44 kDa subunit; General transcription factor IIH polypeptide 2;TFIIH basal transcription factor complex p44 subunit
Observed Band	62kD
Calculated Molecular Weight	62kD
Cell Pathway	Nucleus .
Tissue Specificity	Widely expressed, with higher expression in skeletal muscle.




Function	alternative products:A number of isoforms may be produced. The isoforms may be also produced by incomplete gene duplication,function:Component of the core-TFIIF basal transcription factor involved in nucleotide excision repair (NER) of DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II.,function:Component of the core-TFIIF basal transcription factor involved in nucleotide excision repair (NER) of DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II. The N-terminus interacts with and regulates XPD whereas an intact C-terminus is required for a successful escape of RNAP II from the promoter.,similarity:Belongs to the GTF2H2 family.,similarity:Contains 1 VWFA domain.,subunit:One of the six subunits forming the core-TFIIF basal transcription factor. Interacts with XPB, XPD, GTF2H1 and GTF2H3.,tissue specificity:Widely expressed, with higher express
Background	This gene is part of a 500 kb inverted duplication on chromosome 5q13. This duplicated region contains at least four genes and repetitive elements which make it prone to rearrangements and deletions. The repetitiveness and complexity of the sequence have also caused difficulty in determining the organization of this genomic region. This gene is within the telomeric copy of the duplication. Deletion of this gene sometimes accompanies deletion of the neighboring SMN1 gene in spinal muscular atrophy (SMA) patients but it is unclear if deletion of this gene contributes to the SMA phenotype. This gene encodes the 44 kDa subunit of RNA polymerase II transcription initiation factor IIF which is involved in basal transcription and nucleotide excision repair. Transcript variants for this gene have been described, but their full length nature has not been determined. A second copy of t
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western Blot analysis of COLO205 cells using TFIIH p44 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Western blot analysis of lysates from COLO205 cells, using TF2H2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using TF2H2 antibody.

Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).