



TRAP240 Polyclonal Antibody

Catalog No	YP-Ab-02136
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;IHC;IF;ELISA
Gene Name	MED13L
Immunogen	The antiserum was produced against synthesized peptide derived from human MED13L. AA range:449-498
Specificity	TRAP240 Polyclonal Antibody detects endogenous levels of TRAP240 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	KIAA1025;Mediator complex subunit 13-like;Mediator of RNA polymerase II transcription subunit 13-like;PROSIT240;THRAP2;TRAP240;MED13L;Thyroid hormone receptor-associated protein 2;Thyroid hormone receptor-associated protein complex 240 kDa component-like
Observed Band	250kD
Calculated Molecular Weight	250kD
Cell Pathway	Nucleus .
Tissue Specificity	Highly expressed in brain (cerebellum), heart (aorta), skeletal muscle, kidney, placenta and peripheral blood leukocytes. Highly expressed in fetal brain.
Function	disease:A chromosomal aberration involving MED13L is found in a patient with transposition of the great arteries, dextro-looped and mental retardation. Translocation t(12;17)(q24.1;q21).,disease:Defects in MED13L are a cause of transposition of the great arteries, dextro-looped (DTGA) [MIM:608808]. DTGA consists of complete inversion of the great vessels, so that the aorta incorrectly arises from the right ventricle and the pulmonary artery incorrectly arises from the left ventricle. This creates completely separate pulmonary and systemic circulatory systems, an arrangement that is incompatible with life. Patients often



have atrial and/or ventricular septal defects or other types of shunting that allow some mixing between the circulations in order to support life minimally, but surgical intervention is always required.,function:Component of the Mediator complex, a coactivator involved in

Background

The protein encoded by this gene is a subunit of the Mediator complex, a large complex of proteins that functions as a transcriptional coactivator for most RNA polymerase II-transcribed genes. The encoded protein is involved in early development of the heart and brain. Defects in this gene are a cause of transposition of the great arteries, dextro-looped (DTGA).[provided by RefSeq, Jul 2010],



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using MED13L Antibody. The picture on the right is blocked with the synthesized peptide.