



Neuralized-1 Polyclonal Antibody


Catalog No	YP-Ab-02715
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	NEURL
Immunogen	The antiserum was produced against synthesized peptide derived from human NEURL1. AA range:219-268
Specificity	Neuralized-1 Polyclonal Antibody detects endogenous levels of Neuralized-1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	neu;h-neuralized 1;h neuralized 1;h neu;EC:2.3.2.27;Neuralized-1;NEURL; Neuralized-like protein 1A;NEURL1;NEURL1A;RNF67;h-neu;RING finger protein 67
Observed Band	62kD
Calculated Molecular Weight	62kD
Cell Pathway	Cytoplasm, perinuclear region . Cell membrane ; Peripheral membrane protein . Perikaryon . Cell projection, dendrite . Cell junction, synapse, postsynaptic density . Localized in the cell bodies of the pyramidal neurons and distributed along their apical dendrites. Colocalized with PSD95 in postsynaptic sites. Colocalized with CPEB3 at apical dendrites of CA1 neurons (By similarity). Colocalized with JAG1 at the cell surface. .
Tissue Specificity	Expressed in brain, testis, pituitary gland, pancreas and bone marrow. Also poorly expressed in malignant astrocytomas and several neuroectodermal tumor cell lines. Weakly expressed in medulloblastoma (MB) compared with normal cerebellar tissues.




Function	microtubule cytoskeleton organization, microtubule bundle formation, reproductive developmental process, cell motion, cytoskeleton organization, microtubule-based process, gamete generation, germ cell development, spermatogenesis, spermatid development, sperm axoneme assembly, lactation, flagellum assembly, cellular component assembly involved in morphogenesis, sexual reproduction, cell projection organization, cell projection assembly, sperm motility, mammary gland development, multicellular organism reproduction, cellular component morphogenesis, axoneme assembly, flagellar axoneme assembly, flagellum organization, male gamete generation, spermatid differentiation, reproductive process in a multicellular organism, reproductive cellular process, gland development, cell motility, localization of cell,
Background	NEURL1B (neuralized homolog 1B), also known as neur2 or NEURL3, is a 555 amino acid protein that functions as an E3 ubiquitin-protein ligase. Involved in protein ubiquitination, NEURL1B contains two NHR (neuralized homology repeat) domains and one RING-type zinc finger. NEURL4 (neuralized homolog 4 (Drosophila)), also known as neuralized-like protein 4, is a 1,562 amino acid protein that is highly expressed in a wide variety of tissues. Containing six NHR (neuralized homology repeat) domains, NEURL4 exists as two alternatively spliced isoforms that are encoded by a gene that maps to human chromosome 17p13.1.
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.



Western blot analysis of lysates from MCF-7 cells, using NEURL1 Antibody. The lane on the right is blocked with the synthesized peptide.