



SOD-1 Polyclonal Antibody

Catalog No	YP-Ab-02787
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	SOD1
Immunogen	The antiserum was produced against synthesized peptide derived from human SOD-1. AA range:36-85
Specificity	SOD-1 Polyclonal Antibody detects endogenous levels of SOD-1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ALS1;EC:1.15.1.1;homodimer;hSod1;IPOA;SOD-1;SOD1;Superoxide dismutase [Cu-Zn];Superoxide dismutase 1
Observed Band	18kD
Calculated Molecular Weight	18kD
Cell Pathway	Cytoplasm . Mitochondrion . Nucleus . Predominantly cytoplasmic; the pathogenic variants ALS1 Arg-86 and Ala-94 gradually aggregates and accumulates in mitochondria. .
Tissue Specificity	Colon,Fetal brain cortex,Placenta,
Function	catalytic activity:2 superoxide + 2 H(+) = O(2) + H(2)O(2).,cofactor: Binds 1 copper ion per subunit.,cofactor: Binds 1 zinc ion per subunit.,disease: Defects in SOD1 are the cause of amyotrophic lateral sclerosis type 1 (ALS1) [MIM:105400] . ALS1 is a familial form of amyotrophic lateral sclerosis, a neurodegenerative disorder affecting upper and lower motor neurons and resulting in fatal paralysis. Sensory abnormalities are absent. Death usually occurs within 2 to 5 years. The etiology of amyotrophic lateral sclerosis is likely to be multifactorial, involving both genetic and environmental factors. The disease is inherited in 5-10% of cases leading to familial forms.,function: Destroys radicals which are normally



produced within the cells and which are toxic to biological systems.,
miscellaneous:The protein (both wild-type and ALS1 variants) has a tendency to
form fibrillar aggregates in the

Background

The protein encoded by this gene binds copper and zinc ions and is one of two isozymes responsible for destroying free superoxide radicals in the body. The encoded isozyme is a soluble cytoplasmic protein, acting as a homodimer to convert naturally-occurring but harmful superoxide radicals to molecular oxygen and hydrogen peroxide. The other isozyme is a mitochondrial protein. Mutations in this gene have been implicated as causes of familial amyotrophic lateral sclerosis. Rare transcript variants have been reported for this gene. [provided by RefSeq, Jul 2008],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Chen, Qiong, et al. "Celastrol Alleviates Chronic Obstructive Pulmonary Disease by Inhibiting Cellular Inflammation Induced by Cigarette Smoke via the Ednrb/Kng1 Signaling Pathway." *Frontiers in pharmacology* 9 (2018): 1276.



Western Blot analysis of various cells using SOD-1 Polyclonal Antibody diluted at 1:1000



Western blot analysis of lysate from Jurkat cells, using SOD-1 antibody.