



UBTD2 Polyclonal Antibody


Catalog No	YP-Ab-02821
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	UBTD2
Immunogen	The antiserum was produced against synthesized peptide derived from human UBTD2. AA range:181-230
Specificity	UBTD2 Polyclonal Antibody detects endogenous levels of UBTD2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DCUBP;DC-UbP;ubiquitin domain containing 2;Ubiquitin like protein SB72; UBTD2;Ubiquitin domain-containing protein 2;SB72;Dendritic cell-derived ubiquitin-like protein;Ubiquitin-like protein SB72
Observed Band	26kD
Calculated Molecular Weight	26kD
Cell Pathway	Cytoplasm .
Tissue Specificity	Detected in dendritic cells. Highly expressed in tumor cell lines, but not detectable in most tissues.
Function	similarity:Contains 1 ubiquitin-like domain.,tissue specificity:Detected in dendritic cells. Highly expressed in tumor cell lines, but not detectable in most tissues.,
Background	similarity:Contains 1 ubiquitin-like domain.,tissue specificity:Detected in dendritic cells. Highly expressed in tumor cell lines, but not detectable in most tissues.,
matters needing attention	Avoid repeated freezing and thawing!




Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of UBTD2 Antibody. The lane on the right is blocked with the UBTD2 peptide.



Western blot analysis of the lysates from HUVEC cells using UBTD2 antibody.