



ALDH1A1 Polyclonal Antibody

Catalog No	YP-Ab-02857
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	ALDH1A1
Immunogen	The antiserum was produced against synthesized peptide derived from the N-terminal region of human ALDH1A1. AA range:21-70
Specificity	ALDH1A1 Polyclonal Antibody detects endogenous levels of ALDH1A1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	3-deoxyglucosone dehydrogenase;ALDC;Aldehyde dehydrogenase 1A1;ALDH 1A1;ALDH1 A1;ALDH1A1;Retinal dehydrogenase 1;ALDH1;PUMB1;RALDH 1;RaIDH1;ALDH-E1;ALHDII;Aldehyde dehydrogenase family 1 member A1; Aldehyde dehydrogenase;cytosolic
Observed Band	55kD
Calculated Molecular Weight	55kD
Cell Pathway	Cytoplasm, cytosol . Cell projection, axon .
Tissue Specificity	Expressed by erythrocytes (at protein level).
Function	catalytic activity:Retinal + NAD(+) + H(2)O = retinoate + NADH.,function:Binds free retinal and cellular retinol-binding protein-bound retinal. Can convert/oxidize retinaldehyde to retinoic acid.,pathway:Cofactor metabolism; retinol metabolism., similarity:Belongs to the aldehyde dehydrogenase family., subunit:Homotetramer.,
Background	The protein encoded by this gene belongs to the aldehyde dehydrogenase family. Aldehyde dehydrogenase is the next enzyme after alcohol dehydrogenase in the major pathway of alcohol metabolism. There are two

major aldehyde dehydrogenase isozymes in the liver, cytosolic and mitochondrial, which are encoded by distinct genes, and can be distinguished by their electrophoretic mobility, kinetic properties, and subcellular localization. This gene encodes the cytosolic isozyme. Studies in mice show that through its role in retinol metabolism, this gene may also be involved in the regulation of the metabolic responses to high-fat diet. [provided by RefSeq, Mar 2011],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images




Western Blot analysis of mouse lung, mouse liver cells using ALDH1A1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Western blot analysis of lysate from mouse lung, using ALDH1A1 Antibody.