



UBA1 Polyclonal Antibody

Catalog No	YP-Ab-02874
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	UBA1
Immunogen	The antiserum was produced against synthesized peptide derived from the N-terminal region of human UBA1. AA range:91-140
Specificity	UBA1 Polyclonal Antibody detects endogenous levels of UBA1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	UBA1;GXP1;EC:6.2.1.45;AMCX1;A1ST;Ubiquitin-like modifier-activating enzyme 1;A1S9T;UBE1;Protein A1S9;Ubiquitin-activating enzyme E1
Observed Band	118kD
Calculated Molecular Weight	118kD
Cell Pathway	Cytoplasm . Mitochondrion . Nucleus .; [Isoform 1]: Nucleus .; [Isoform 2]: Cytoplasm .
Tissue Specificity	Detected in erythrocytes (at protein level). Ubiquitous.
Function	disease:Defects in UBA1 are the cause of spinal muscular atrophy X-linked type 2 (SMAX2) [MIM:301830]; also known as X-linked lethal infantile spinal muscular atrophy, distal X-linked arthrogryposis multiplex congenita or X-linked arthrogryposis type 1 (AMCX1). Spinal muscular atrophy refers to a group of neuromuscular disorders characterized by degeneration of the anterior horn cells of the spinal cord, leading to symmetrical muscle weakness and atrophy. SMAX2 is a lethal infantile form presenting with hypotonia, areflexia, and multiple congenital contractures.,function:Activates ubiquitin by first adenylating its C-terminal glycine residue with ATP, and thereafter linking this residue to the side chain of a cysteine residue in E1, yielding an ubiquitin-E1 thioester and free



AMP.,miscellaneous:There are two active sites within the E1 molecule, allowing it to accommodate two ubiquitin mo

Background

The protein encoded by this gene catalyzes the first step in ubiquitin conjugation to mark cellular proteins for degradation. This gene complements an X-linked mouse temperature-sensitive defect in DNA synthesis, and thus may function in DNA repair. It is part of a gene cluster on chromosome Xp11.23. Alternatively spliced transcript variants that encode the same protein have been described. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of PC12 cells using UBA1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000