



Mox1 Polyclonal Antibody

Catalog No	YP-Ab-02895
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	NOX1 MOX1 NOH1
Immunogen	Synthetic peptide from human protein at AA range: 210-260
Specificity	The antibody detects endogenous Mox1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000,IHC-p 1:500-200, ELISA 1:10000-20000. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EC:1.6.3.-;GP91 2;Mitogenic oxidase 1;MOX 1;MOX1;NOX1 MOX1 NOH1; NADPH oxidase 1 (NOX-1) (EC 1.-.-.-) (Mitogenic oxidase 1) (MOX-1) (NADH/ NADPH mitogenic oxidase subunit P65-MOX) (NOH-1);NADPH oxidase 1 (NOX-1;EC 1.-.-.-;MOX-1;NADH/NADPH mitogenic oxidase subunit P65-MOX;NOH-1)
Observed Band	65kD
Calculated Molecular Weight	65kD
Cell Pathway	Cell projection, invadopodium membrane ; Multi-pass membrane protein . Cell membrane .
Tissue Specificity	NOH-1L is detected in colon, uterus, prostate, and colon carcinoma, but not in peripheral blood leukocytes. NOH-1S is detected only in colon and colon carcinoma cells.
Function	cofactor:FAD .,cofactor:NADP .,enzyme regulation:The oxidase activity is potentiated by NOXA1 and NOXO1.,function:NOH-1S is a voltage-gated proton channel that mediates the H(+) currents of resting phagocytes and other tissues. It participates in the regulation of cellular pH and is blocked by zinc. NOH-1L is a pyridine nucleotide-dependent oxidoreductase that generates superoxide and might conduct H(+) ions as part of its electron transport mechanism, whereas NOH-1S does not contain an electron transport chain.,similarity:Contains 1 FAD-



binding FR-type domain.,similarity:Contains 1 ferric oxidoreductase domain.,
subunit:NOX1, NOXA1, NOXO1, RAC1 and CYBA forms a functional multimeric
complex supporting ROS production. Interacts with NOXA1 and NOXO1.,tissue
specificity:NOH-1L is detected in colon, uterus, prostate, and colon carcinoma,
but not in peripheral blood leukocytes. NOH-1S is dete

Background

This gene encodes a member of the NADPH oxidase family of enzymes responsible for the catalytic one-electron transfer of oxygen to generate superoxide or hydrogen peroxide. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Nov 2012],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images




Western blot analysis of 293T Hela VEC KB mouse-kidney 3T3 lysate, antibody was diluted at 2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-stomach, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-stomach, antibody was diluted at 1:200