



S100B mouse mAb

Catalog No	YP-Ab-02965
Isotype	IgG
Reactivity	Rat
Applications	WB
Gene Name	S100B
Immunogen	Human S100B synthetic peptide conjugated to KLH.
Specificity	This antibody detects endogenous levels of S100B and does not cross-react with related proteins.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Dilution	wb dilution 1:500
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	S100 beta;Protein S100 B;Protein S100-B;S100 calcium-binding protein B;S-100 protein beta chain;S100B;NEF;S 100 calcium binding protein beta chain;S 100 protein beta chain;S-100 protein subunit beta;S100;S100 calcium binding protein beta (neural);S100 protein beta chain;S100B_HUMAN;S100beta.
Observed Band	10kD
Calculated Molecular Weight	10kD
Cell Pathway	Cytoplasm . Nucleus .
Tissue Specificity	Although predominant among the water-soluble brain proteins, S100 is also found in a variety of other tissues.
Function	function:Weakly binds calcium but binds zinc very tightly-distinct binding sites with different affinities exist for both ions on each monomer. Physiological concentrations of potassium ion antagonize the binding of both divalent cations, especially affecting high-affinity calcium-binding sites. Binds to and initiates the activation of STK38 by releasing autoinhibitory intramolecular interactions within the kinase.,miscellaneous:In addition to metal-ion binding, this protein is involved with the regulation of protein phosphorylation in brain tissue.,similarity:Belongs to the S-101 family.,similarity:Contains 2 EF-hand domains.,



subunit:Dimer of either two alpha chains, or two beta chains, or one alpha and one beta chain. The S100B dimer binds two molecules of STK38 (By similarity). The S100B dimer interacts with two molecules of CAPZA1.,tissue specificity:Although predominant among the wa

Background

The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21; however, this gene is located at 21q22.3. This protein may function in Neurite extension, proliferation of melanoma cells, stimulation of Ca²⁺ fluxes, inhibition of PKC-mediated phosphorylation, astrocytosis and axonal proliferation, and inhibition of microtubule assembly. Chromosomal rearrangements and altered expression of this gene have been implicated in several neurological, neoplastic, and other types of diseases, including Alzheimer's disease, Down's syndrome, epilepsy


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of extracts from Rat Brain cell lysates using S100B mouse mAb (1:500 diluted). Predicted band size: 10kDa. Observed band size: 10kDa.