



CIB1 rabbit pAb

Catalog No	YP-Ab-03243
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB; ELISA
Gene Name	CIB1 CIB KIP PRKDCIP
Immunogen	Synthesized peptide derived from human CIB1 AA range: 70-150
Specificity	This antibody detects endogenous levels of Human CIB1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CIB;CIBP;KIP1;PRKDCIP;SIP2-28;CIB1;CIB1 CIB KIP PRKDCIP;Calcium and integrin-binding protein 1 (CIB;Calcium- and integrin-binding protein;Calmyrin; DNA-PKcs-interacting protein;Kinase-interacting protein;KIP;SNK-interacting protein 2-28;SIP2-28)
Observed Band	22kDa
Calculated Molecular Weight	22kDa
Cell Pathway	Membrane; Lipid-anchor. Cell membrane, sarcolemma. Cell membrane. Apical cell membrane. Cell projection, ruffle membrane. Cell projection, filopodium tip. Cell projection, growth cone . Cell projection, lamellipodium . Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, perinuclear region. Nucleus . Cell projection, neuron projection . Perikaryon . Colocalized with PPP3R1 at the cell membrane of cardiomyocytes in the hypertrophic heart (By similarity). Colocalized with NBR1 to the perinuclear region. Colocalizes with TAS1R2 in apical regions of taste receptor cells. Colocalized with RAC3 in the perinuclear area and at the cell periphery. Colocalized with PAK1 within membrane ruffles during cell spreading upon readhesion to fibr



Tissue Specificity	Ubiquitously expressed. Expressed in the epidermis, hair follicles and keratinocytes (PubMed:30068544). Detected in platelets and in cell lines of megakaryocytic and erythrocytic lineages. Both isoform 1 and isoform 2 are detected in various cancer cell lines, with isoform 2 being the predominant form (at protein level).
Function	function:May convert the inactive conformation of integrin alpha-IIb/beta3 to an active form through the binding to the integrin cytoplasmic domain., similarity:Contains 2 EF-hand domains.,subunit:Monomer. Interacts with the heterodimeric integrin alpha-IIb/beta3. Interacts with the protein kinases PLK2/SNK and with the region immediately upstream of the kinase domain of DNA-PK. Interacts with PSEN2. Interacts with MYO1C.,tissue specificity:Ubiquitous.,
Background	This gene encodes a member of the EF-hand domain-containing calcium-binding superfamily. The encoded protein interacts with many other proteins, including the platelet integrin alpha-IIb-beta-3, DNA-dependent protein kinase, presenilin-2, focal adhesion kinase, p21 activated kinase, and protein kinase D. The encoded protein may be involved in cell survival and proliferation, and is associated with several disease states including cancer and Alzheimer's disease. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2013],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images