



AAT Monoclonal Antibody

Catalog No	YP-Ab-03380
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	SERPINA1
Immunogen	Purified recombinant fragment of human AAT expressed in E. Coli.
Specificity	AAT Monoclonal Antibody detects endogenous levels of AAT protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AAT 1;AAT-1;AAT1 alpha;AAT1;AAT1_HUMAN;AAT1alpha;AMY 1 associating protein expressed in testis 1;AMY-1-associating protein expressed in testis 1; Chromosome 3 open reading frame 15;MYCBP binding protein;AAT;SERPINA1; Alpha-1-antitrypsin;PI;Alpha-1 protease inhibitor;Alpha-1-antiproteinase;Serpina1
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Secreted. Endoplasmic reticulum. The S and Z allele are not secreted effectively and accumulate intracellularly in the endoplasmic reticulum.; [Short peptide from AAT]: Secreted, extracellular space, extracellular matrix.
Tissue Specificity	Ubiquitous. Expressed in leukocytes and plasma.
Function	disease:Deficiency of the normal inhibitor in individuals homozygous for allele Z or M-Malton can result in the development of chronic emphysema or infantile liver cirrhosis.,disease:The major physiological function of AAT is the protection of the lower respiratory tract against proteolytic destruction by human leukocyte elastase (HLE). A hereditary deficiency of AAT, is associated with a 20-30 fold increased risk of developing chronic obstructive pulmonary disease., disease:Variant Pittsburgh is the cause of bleeding diathesis.,domain:The



reactive center loop (RCL) extends out from the body of the protein and directs binding to the target protease. The protease cleaves the serpin at the reactive site within the RCL, establishing a covalent linkage between the carboxyl group of the serpin reactive site and the serine hydroxyl of the protease. The resulting inactive serpin-protease complex

Background

The protein encoded by this gene is secreted and is a serine protease inhibitor whose targets include elastase, plasmin, thrombin, trypsin, chymotrypsin, and plasminogen activator. Defects in this gene can cause emphysema or liver disease. Several transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis using AAT Monoclonal Antibody against human plasma (1) and NIH/3T3 cell lysate (2).