



Fibrinogen β Monoclonal Antibody

Catalog No	YP-Ab-03396
Isotype	IgG
Reactivity	Human
Applications	WB;ELISA
Gene Name	FGB
Immunogen	Purified recombinant fragment of human Fibrinogen β (aa30-300) expressed in E. Coli.
Specificity	Fibrinogen β Monoclonal Antibody detects endogenous levels of Fibrinogen β protein.
Formulation	Ascitic fluid containing 0.03% sodium azide, 0.5% BSA, 50% glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	Fibrinogen β ; FGB; Fibrinogen beta chain
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Secreted .
Tissue Specificity	Detected in blood plasma (at protein level).
Function	disease: Defects in FGB are a cause of congenital afibrinogenemia [MIM:202400] . This rare autosomal recessive disorder is characterized by bleeding that varies from mild to severe and by complete absence or extremely low levels of plasma and platelet fibrinogen., disease: Defects in FGB are a cause of thrombophilia., domain: A long coiled coil structure formed by 3 polypeptide chains connects the central nodule to the C-terminal domains (distal nodules). The long C-terminal ends of the alpha chains fold back, contributing a fourth strand to the coiled coil structure., function: Fibrinogen has a double function: yielding monomers that polymerize into fibrin and acting as a cofactor in platelet aggregation., online information: Fibrinogen entry, online information: The Singapore human mutation and polymorphism database, PTM: Conversion of fibrinogen to fibrin is triggered



by thrombin, which cleaves fib

Background

The protein encoded by this gene is the beta component of fibrinogen, a blood-borne glycoprotein comprised of three pairs of nonidentical polypeptide chains. Following vascular injury, fibrinogen is cleaved by thrombin to form fibrin which is the most abundant component of blood clots. In addition, various cleavage products of fibrinogen and fibrin regulate cell adhesion and spreading, display vasoconstrictor and chemotactic activities, and are mitogens for several cell types. Mutations in this gene lead to several disorders, including afibrinogenemia, dysfibrinogenemia, hypodysfibrinogenemia and thrombotic tendency. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jun 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis using Fibrinogen β Monoclonal Antibody against human plasma (1).