



IMP-3 Monoclonal Antibody

Catalog No	YP-Ab-03408
Isotype	IgG
Reactivity	Human
Applications	WB;IHC;IF;ELISA
Gene Name	IGF2BP3
Immunogen	Purified recombinant fragment of human IMP-3 expressed in E. Coli.
Specificity	IMP-3 Monoclonal Antibody detects endogenous levels of IMP-3 protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	WB: 1/500 - 1/2000. IHC: 1/200 - 1/1000. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	BRMS2;C15orf12;Chromosome 15 open reading frame 12;DKFZp586L118; FLJ1968;imp3;U3 small nucleolar ribonucleoprotein;homolog (yeast); IMP3_HUMAN;Mitochondrial ribosomal protein S4;MRPS4;OTTHUMP183524; U3 small nucleolar ribonucleoprotein protein imp3;U3 snoRNP protein 3 homolog;U3 snoRNP protein imp3;IMP-3;IGF2BP3;Insulin-like growth factor 2 mRNA-binding protein 3;KOC1;VICKZ3;IGF2 mRNA-binding protein 3;IGF-II mRNA-binding protein 3;KH domain-containing protein overexpressed in cancer; hKOC;VICKZ family member 3
Observed Band	64kD
Calculated Molecular Weight	64kD
Cell Pathway	Nucleus. Cytoplasm. Cytoplasm, P-body . Cytoplasm, Stress granule . Found in lamellipodia of the leading edge, in the perinuclear region, and beneath the plasma membrane. The subcytoplasmic localization is cell specific and regulated by cell contact and growth. Localized at the connecting piece and the tail of the spermatozoa. Colocalized with CD44 mRNA in RNP granules. In response to cellular stress, such as oxidative stress, recruited to stress granules.
Tissue Specificity	Expressed in fetal liver, fetal lung, fetal kidney, fetal thymus, fetal placenta, fetal follicles of ovary and gonocytes of testis, growing oocytes, spermatogonia and semen (at protein level). Expressed in cervix adenocarcinoma, in testicular, pancreatic and renal-cell carcinomas (at protein level). Expressed ubiquitously



during fetal development at 8 and 14 weeks of gestation. Expressed in ovary, testis, brain, placenta, pancreatic cancer tissues and pancreatic cancer cell lines.

Function

disease: Autoantibodies against IGF2BP3 are detected in sera from some patients with a variety of carcinomas., function: RNA-binding protein that act as a regulator of mRNA translation and stability. Binds to the 5'-UTR of the insulin-like growth factor 2 (IGF2) mRNAs. Binds to sequences in the 3'-UTR of CD44 mRNA., similarity: Belongs to the RRM IMP/VICKZ family., similarity: Contains 2 RRM (RNA recognition motif) domains., similarity: Contains 4 KH domains., subcellular location: Found in lamellipodia of the leading edge, in the perinuclear region, and beneath the plasma membrane. The subcytoplasmic localization is cell specific and regulated by cell contact and growth. Localized at the connecting piece and the tail of the spermatozoa. Colocalized with CD44 mRNA in RNP granules., subunit: Homodimer and multimer., tissue specificity: Expressed in fetal liver, fetal lung, fetal kidney, fetal thymus, fe

Background

The protein encoded by this gene is primarily found in the nucleolus, where it can bind to the 5' UTR of the insulin-like growth factor II leader 3 mRNA and may repress translation of insulin-like growth factor II during late development. The encoded protein contains several KH domains, which are important in RNA binding and are known to be involved in RNA synthesis and metabolism. A pseudogene exists on chromosome 7, and there are putative pseudogenes on other chromosomes. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis using IMP-3 Monoclonal Antibody against Jurkat (1), K562 (2) and NTERA-2 (3) cell lysate.



Immunohistochemistry analysis of paraffin-embedded lung cancer (left) and colon tumour tissues (right) with DAB staining using IMP-3 Monoclonal Antibody.