



SIRT1 (phospho Ser47) Polyclonal Antibody

Catalog No	YP-Ab-03547
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA;IHC
Gene Name	SIRT1
Immunogen	Synthesized phospho-peptide around the phosphorylation site of human SIRT1 (phospho Ser47)
Specificity	Phospho-SIRT1 (S47) Polyclonal Antibody detects endogenous levels of SIRT1 protein only when phosphorylated at S47.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	75SirT1;hSIR2;hSIRT1;HST2;S. cerevisiae;homolog of;NAD dependent deacetylase sirtuin 1;NAD dependent protein deacetylase sirtuin 1; OTTHUMP198111;OTTHUMP198112;Regulatory protein SIR2 homolog 1; SIR1_HUMAN;SIR2;SIR2 like 1;SIR2 like protein 1;S.cerevisiae;homolog-like 1; SIR2-like protein 1;SIR2ALPHA;SIR2L1;Sirt1;SirtT1 75 kDa fragment;Sirtuin (silent mating type information regulation 2 homolog) 1 (S. cerevisiae);Sirtuin 1; Sirtuin type 1;NAD-dependent protein deacetylase sirtuin-1
Observed Band	85-110kD
Calculated Molecular Weight	85-110kD
Cell Pathway	Nucleus, PML body . Cytoplasm . Nucleus . Recruited to the nuclear bodies via its interaction with PML (PubMed:12006491). Colocalized with APEX1 in the nucleus (PubMed:19934257). May be found in nucleolus, nuclear euchromatin, heterochromatin and inner membrane (PubMed:15469825). Shuttles between nucleus and cytoplasm (By similarity). Colocalizes in the nucleus with XBP1 isoform 2 (PubMed:20955178). .; [SirtT1 75 kDa fragment]: Cytoplasm . Mitochondrion .
Tissue Specificity	Widely expressed.



Function

catalytic activity: NAD(+) + an acetylprotein = nicotinamide + O-acetyl-ADP-ribose + a protein., cofactor: Binds 1 zinc ion per subunit., enzyme regulation: Inhibited by nicotinamide. Activated by resveratrol (3,5,4'-trihydroxy-trans-stilbene), butein (3,4,2',4'-tetrahydroxychalcone), piceatannol (3,5,3',4'-tetrahydroxy-trans-stilbene), Isoliquiritigenin (4,2',4'-trihydroxychalcone), fisetin (3,7,3',4'-tetrahydroxyflavone) and quercetin (3,5,7,3',4'-pentahydroxyflavone). RPS19BP1/AROS acts as a positive regulator of deacetylation activity., function: NAD-dependent deacetylase, which regulates processes such as apoptosis and muscle differentiation by deacetylating key proteins. Deacetylates 'Lys-382' of p53/TP53 and impairs its ability to induce proapoptotic program and modulate cell senescence. Deacetylates TAF1B and thereby represses rDNA transcription by the RNA polymerase I. Involved in HES1

Background

This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).