



CEP57 Polyclonal Antibody

Catalog No	YP-Ab-03763
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	CEP57
Immunogen	The antiserum was produced against synthesized peptide derived from human CEP57. AA range:241-290
Specificity	CEP57 Polyclonal Antibody detects endogenous levels of CEP57 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	centrosomal protein 57kDa;Centrosomal protein of 57 kDa;FGF2 interacting protein;KIAA92;PIG8;CEP57;KIAA0092;TSP57;FGF2-interacting protein;Testis-specific protein 57;Translokin
Observed Band	50kD
Calculated Molecular Weight	50kD
Cell Pathway	Nucleus . Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .
Tissue Specificity	Ubiquitous.
Function	function:Mediates nuclear translocation and mitogenic activity of the internalized growth factor FGF2.,similarity:Belongs to the translokin family.,subcellular location:Associates with microtubules and the centrosome.,subunit:Homodimer. Interacts with FGF2 and RAP80. Does not interact with FGF1 or FGF2 isoform 24 kDa.,tissue specificity:Ubiquitous.,
Background	This gene encodes a cytoplasmic protein called Translokin. This protein localizes to the centrosome and has a function in microtubular stabilization. The N-terminal half of this protein is required for its centrosome localization and for



its multimerization, and the C-terminal half is required for nucleating, bundling and anchoring microtubules to the centrosomes. This protein specifically interacts with fibroblast growth factor 2 (FGF2), sorting nexin 6, Ran-binding protein M and the kinesins KIF3A and KIF3B, and thus mediates the nuclear translocation and mitogenic activity of the FGF2. It also interacts with cyclin D1 and controls nucleocytoplasmic distribution of the cyclin D1 in quiescent cells. This protein is crucial for maintaining correct chromosomal number during cell division. Mutations in this gene cause mosaic variegated aneuploidy syndrome, a rare autosomal recessive disorder. Multiple


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from COLO and LOVO cells, using CEP57 Antibody. The lane on the right is blocked with the synthesized peptide.