



CES2 Polyclonal Antibody

Catalog No	YP-Ab-03769
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	CES2
Immunogen	Synthesized peptide derived from the Internal region of human CES2.
Specificity	CES2 Polyclonal Antibody detects endogenous levels of CES2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Carboxylesterase 2;CE 2;CES2;CES2A1;hCE 2;iCE;PCE 2;Cocaine esterase; CE-2;hCE-2;Methylumbelliferyl-acetate deacetylase 2
Observed Band	61kD
Calculated Molecular Weight	61kD
Cell Pathway	Endoplasmic reticulum lumen .
Tissue Specificity	Preferentially expressed in intestine with moderate expression in liver. Within the intestine, highest expression is found in small intestine with lower expression in colon and rectum.
Function	catalytic activity:A carboxylic ester + H(2)O = an alcohol + a carboxylate., function:Involved in the detoxification of xenobiotics and in the activation of ester and amide prodrugs. Shows high catalytic efficiency for hydrolysis of 4-methumbelliferyl acetate, heroin and 6-monoacetylmorphine., PTM:Glycosylated.,similarity:Belongs to the type-B carboxylesterase/lipase family.,subunit:Monomer.,tissue specificity:Preferentially expressed in intestine with moderate expression in liver. Within the intestine, highest expression is found in small intestine with lower expression in colon and rectum.,



Background

This gene encodes a member of the carboxylesterase large family. The family members are responsible for the hydrolysis or transesterification of various xenobiotics, such as cocaine and heroin, and endogenous substrates with ester, thioester, or amide bonds. They may participate in fatty acyl and cholesterol ester metabolism, and may play a role in the blood-brain barrier system. The protein encoded by this gene is the major intestinal enzyme and functions in intestine drug clearance. Alternatively spliced transcript variants have been found for this gene.[provided by RefSeq, Oct 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).