



CHST13 Polyclonal Antibody

Catalog No	YP-Ab-03771
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	CHST13
Immunogen	The antiserum was produced against synthesized peptide derived from human CHST13. AA range:211-260
Specificity	CHST13 Polyclonal Antibody detects endogenous levels of CHST13 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	C4ST 3;C4ST-3;C4ST3;Carbohydrate (chondroitin 4) sulfotransferase 13; Carbohydrate sulfotransferase 13;Chondroitin 4 O sulfotransferase 3; Chondroitin 4 sulfotransferase 3;Chondroitin 4-O-sulfotransferase 3;Chondroitin 4-sulfotransferase 3;CHST 13;CHST13;CHSTD_HUMAN;MGC119278; MGC119279;MGC119281
Observed Band	39kD
Calculated Molecular Weight	39kD
Cell Pathway	Golgi apparatus membrane ; Single-pass type II membrane protein .
Tissue Specificity	Highly expressed in adult liver. Expressed at lower level in kidney, lymph nodes and fetal kidney.
Function	catalytic activity:3'-phosphoadenylyl sulfate + chondroitin = adenosine 3',5'-bisphosphate + chondroitin 4'-sulfate.;function:Catalyzes the transfer of sulfate to position 4 of the N-acetylgalactosamine (GalNAc) residue of chondroitin. Chondroitin sulfate constitutes the predominant proteoglycan present in cartilage and is distributed on the surfaces of many cells and extracellular matrices. Transfers sulfate to the C4 hydroxyl of beta1,4-linked GalNAc that is substituted with a beta-linked glucuronic acid at the C-3 hydroxyl. No activity toward



dermatan.,similarity:Belongs to the sulfotransferase 2 family.,tissue
specificity:Highly expressed in adult liver. Expressed at lower level in kidney,
lymph nodes and fetal kidney.,

Background

The protein encoded by this gene belongs to the sulfotransferase 2 family. It is localized to the golgi membrane, and catalyzes the transfer of sulfate to the C4 hydroxyl of beta-1,4-linked N-acetylgalactosamine (GalNAc) flanked by glucuronic acid residue in chondroitin. Chondroitin sulfate constitutes the predominant proteoglycan present in cartilage and is distributed on the surfaces of many cells and extracellular matrices. [provided by RefSeq, Aug 2011],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western Blot analysis of K562 cells using CHST13 Polyclonal Antibody



Western blot analysis of CHST13 Antibody. The lane on the right is blocked with the CHST13 peptide.



Immunohistochemistry analysis of paraffin-embedded human testis, using CHST13 Antibody. The lane on the right is blocked with the CHST13 peptide.