



eIF3α Polyclonal Antibody

Catalog No	YP-Ab-03844
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	EIF3J
Immunogen	Synthesized peptide derived from eIF3α . at AA range: 40-120
Specificity	eIF3α Polyclonal Antibody detects endogenous levels of eIF3α protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	centrosomin homolog;cytoplasmic protein p167;eIF 3 theta;eIF-3-theta;EIF3; eIF3 p167;eIF3 p17;eIF3 p18;EIF3 p18 subunit;eIF3 p185;eIF3 theta;eIF3a; EIF3A_HUMAN;EIF3S1;eukaryotic translation initiation factor 3 subunit 1 (theta 15 17kD);eukaryotic translation initiation factor 3 subunit 1 (theta 17kD); eukaryotic translation initiation factor 3 subunit 1 17kDa;Eukaryotic translation initiation factor 3 subunit 1;eukaryotic translation initiation factor 3 subunit 1 theta 15 17kDa;Eukaryotic translation initiation factor 3 subunit A; OTTHUMP2589;P167;p18;p185;TIF32;eIF3α;EIF3J;Eukaryotic translation initiation factor 3 subunit J;eIF-3-alpha;eIF3 p35
Observed Band	30kD
Calculated Molecular Weight	30kD
Cell Pathway	Cytoplasm .
Tissue Specificity	Epithelium,Fetal liver,Lung carcinoma,Ovarian carci
Function	function:Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the



43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. This subunit binds directly within the mRNA entry channel of the 40S ribosome to the aminoacyl (A) site. It may regulate the interaction between the 43S PIC and mRNA.,mass spectrometry: PubMed:17322308,mass spectrometry: PubMed:18599441,P

Background

This gene encodes a core subunit of the eukaryotic initiation factor 3 complex, which participates in the initiation of translation by aiding in the recruitment of protein and mRNA components to the 40S ribosome. There are pseudogenes for this gene on chromosomes 1, 3, and 9. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Sep 2013],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images

SH-SY5Y 293T 3T3



Tubulin 55KD

Western Blot analysis of various cells using Antibody diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



eIF3α 30KD