



Factor VIII Polyclonal Antibody

Catalog No	YP-Ab-03865
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;IHC;IF;ELISA
Gene Name	F8
Immunogen	The antiserum was produced against synthesized peptide derived from human Factor VIII. AA range:2161-2210
Specificity	Factor VIII Polyclonal Antibody detects endogenous levels of Factor VIII protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AHF;Antihemophilic factor;Coagulation factor VIII;procoagulant component; coagulation factor VIIIc;DXS1253E;F8;F8b;F8c;FA8_HUMAN;factor VIII F8B; Factor VIIIa light chain;FactorVIII;FVIII;Hema;Hemophilia A;Hemophilia;classic; OTTHUMP61446;OTTHUMP196174;Factor VIII
Observed Band	300kD
Calculated Molecular Weight	300kD
Cell Pathway	Secreted, extracellular space.
Tissue Specificity	Brain,Hippocampus,Kidney,Plasma,
Function	disease:Defects in F8 are the cause of hemophilia A (HEMA) [MIM:306700]. HEMA is a common recessive X-linked coagulation disorder. The frequency of hemophilia A is 1-2 in 10,000 male births in all ethnic groups. About 50% of patients have severe hemophilia A with F8C activity less than 1% of normal; they have frequent spontaneous bleeding into joints, muscles and internal organs. Moderately severe hemophilia A occurs in about 10% of patients; F8C activity is 2-5% of normal, and there is bleeding after minor trauma. Mild hemophilia A, which occurs in 30-40% of patients, is associated with F8C activity of 5-30% and bleeding occurs only after significant trauma or surgery. Of



particular interest for the understanding of the function of F8C is the category of CRM (cross-reacting material) positive patients (approximately 5%) that have considerable amount of F8C in their plasma (at least 30%)

Background

This gene encodes coagulation factor VIII, which participates in the intrinsic pathway of blood coagulation; factor VIII is a cofactor for factor IXa which, in the presence of Ca^{+2} and phospholipids, converts factor X to the activated form Xa. This gene produces two alternatively spliced transcripts. Transcript variant 1 encodes a large glycoprotein, isoform a, which circulates in plasma and associates with von Willebrand factor in a noncovalent complex. This protein undergoes multiple cleavage events. Transcript variant 2 encodes a putative small protein, isoform b, which consists primarily of the phospholipid binding domain of factor VIIIc. This binding domain is essential for coagulant activity. Defects in this gene results in hemophilia A, a common recessive X-linked coagulation disorder. [provided by RefSeq, Jul 2008],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images





Western Blot analysis of various cells using Factor VIII Polyclonal Antibody



Western Blot analysis of HuvEc cells using Factor VIII Polyclonal Antibody



Western blot analysis of lysates from HUVEC cells, using Factor VIII Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 30min).