



FucT-III Polyclonal Antibody

Catalog No	YP-Ab-03880
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	FUT3
Immunogen	The antiserum was produced against synthesized peptide derived from human FUT3. AA range:91-140
Specificity	FucT-III Polyclonal Antibody detects endogenous levels of FucT-III protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	LE;Les;FT3B;CD174;FucT-III;FUT3;Galactoside 3(4)-L-fucosyltransferase;Blood group Lewis alpha-4-fucosyltransferase;Lewis FT;Fucosyltransferase 3; Fucosyltransferase III
Observed Band	42kD
Calculated Molecular Weight	42kD
Cell Pathway	Golgi apparatus, Golgi stack membrane; Single-pass type II membrane protein . Membrane-bound form in trans cisternae of Golgi.
Tissue Specificity	Highly expressed in stomach, colon, small intestine, lung and kidney and to a lesser extent in salivary gland, bladder, uterus and liver.
Function	catalytic activity:GDP-beta-L-fucose + beta-D-galactosyl-(1->3)-N-acetyl-D-glucosaminyl-R = GDP + beta-D-galactosyl-(1->3)-(alpha-L-fucosyl-(1->4))-N-acetyl-beta-D-glucosaminyl- R.,function:May catalyze alpha-1,3 and alpha-1,4 glycosidic linkages involved in the expression of Vim-2, Lewis A, Lewis B, sialyl Lewis X and Lewis X/SSEA-1 antigens. May be involved in blood group Lewis determination; Lewis-positive (Le(+)) individuals have an active enzyme while Lewis-negative (Le(-)) individuals have an inactive enzyme.,miscellaneous:Also acts on the corresponding 1,4-galactosyl derivative, forming 1,3-L-fucosyl links., online information:Blood group antigen gene mutation database,online



information:Fucosyltransferase 3,online information:GlycoGene database,
pathway:Protein modification; protein glycosylation.,similarity:Belongs to the
glycosyltransferase 10 family.,subcellular location:Membrane

Background

The Lewis histo-blood group system comprises a set of fucosylated glycosphingolipids that are synthesized by exocrine epithelial cells and circulate in body fluids. The glycosphingolipids function in embryogenesis, tissue differentiation, tumor metastasis, inflammation, and bacterial adhesion. They are secondarily absorbed to red blood cells giving rise to their Lewis phenotype. This gene is a member of the fucosyltransferase family, which catalyzes the addition of fucose to precursor polysaccharides in the last step of Lewis antigen biosynthesis. It encodes an enzyme with alpha(1,3)-fucosyltransferase and alpha(1,4)-fucosyltransferase activities. Mutations in this gene are responsible for the majority of Lewis antigen-negative phenotypes. Multiple alternatively spliced variants, encoding the same protein, have been found for this gene. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HCT116 cells,
primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).