



MRP-L35 Polyclonal Antibody

Catalog No	YP-Ab-04346
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;IHC
Gene Name	MRPL35
Immunogen	Synthesized peptide derived from MRP-L35 . at AA range: 80-160
Specificity	MRP-L35 Polyclonal Antibody detects endogenous levels of MRP-L35 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MRPL 35;MRP L35;L35mt;MRP-L35;MRPL35;39S ribosomal protein L35 mitochondrial;BM-007;39S ribosomal protein L35;mitochondrial
Observed Band	21kD
Calculated Molecular Weight	21kD
Cell Pathway	Mitochondrion .
Tissue Specificity	Bone marrow,Liver,Stomach,
Function	similarity:Belongs to the ribosomal protein L35P family.,
Background	Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 39S subunit protein. Sequence analysis identified three transcript variants. Pseudogenes



corresponding to this gene are found on chromosomes 6p, 10q, and Xp.
[provided by RefSeq, Jul 2008],

**matters needing
attention**


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western blot analysis of 293T COLO using MRP-L35 antibody. Antibody was diluted at 1:2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of 293 using MRP-L35 antibody. Antibody was diluted at 1:2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human uterus. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).