



CHD9 Polyclonal Antibody

Catalog No	YP-Ab-05469
Isotype	IgG
Reactivity	Human;Mouse
Applications	IHC;IF
Gene Name	CHD9 KIAA0308 KISH2 PRIC320 AD-013 x0008
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	CHD9 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC-p 1:50-300. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AD013;AD-013;ATP-dependent helicase CHD9;CHD-9;Chromatin-related mesenchymal modulator;CHD9;CHD9 KIAA0308 KISH2 PRIC320 AD-013 x0008;Chromodomain-helicase-DNA-binding protein 9 (CHD-9) (EC 3.6.4.12) (ATP-dependent helicase CHD9) (Chromatin-related mesenchymal modulator) (CReMM) (Chromatin-remodeling factor CHROM1) (Kismet homolog 2
Observed Band	318kD
Calculated Molecular Weight	318kD
Cell Pathway	Cytoplasm . Nucleus .
Tissue Specificity	Widely expressed at low levels. In bone marrow, expression is restricted to osteoprogenitor cells adjacent to mature osteoblasts.
Function	function:Acts as a transcriptional coactivator for PPARA and possibly other nuclear receptors. Proposed to be a ATP-dependent chromatin remodeling protein. Has DNA-dependent ATPase activity and binds to A/T-rich DNA. Associates with A/T-rich regulatory regions in promoters of genes that participate in the differentiation of progenitors during osteogenesis., PTM:Phosphorylated on serine and tyrosine residues.,sequence caution:Wrong choice of frame.,similarity:Belongs to the SNF2/RAD54 helicase family.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1



helicase C-terminal domain.,similarity:Contains 2 chromo domains.,
subunit:Interacts with PPARA. Probably interacts with ESR1 and NR1I3.,tissue
specificity:Widely expressed at low levels. In bone marrow, expression is
restricted to osteoprogenitor cells adjacent to mature osteoblasts.,

Background

function:Acts as a transcriptional coactivator for PPARA and possibly other
nuclear receptors. Proposed to be a ATP-dependent chromatin remodeling
protein. Has DNA-dependent ATPase activity and binds to A/T-rich DNA.
Associates with A/T-rich regulatory regions in promoters of genes that
participate in the differentiation of progenitors during osteogenesis.,
PTM:Phosphorylated on serine and tyrosine residues.,sequence caution:Wrong
choice of frame.,similarity:Belongs to the SNF2/RAD54 helicase family.,
similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1
helicase C-terminal domain.,similarity:Contains 2 chromo domains.,
subunit:Interacts with PPARA. Probably interacts with ESR1 and NR1I3.,tissue
specificity:Widely expressed at low levels. In bone marrow, expression is
restricted to osteoprogenitor cells adjacent to mature osteoblasts.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For
more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded
human liver cancer. 1, Antibody was diluted at 1:200(4°
overnight). 2, Tris-EDTA,pH9.0 was used for antigen
retrieval. 3,Secondary antibody was diluted at
1:200(room temperature, 45min).