



PHX2B Polyclonal Antibody

Catalog No	YP-Ab-05898
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	PHOX2B PMX2B
Immunogen	Synthesized peptide derived from human protein . at AA range: 140-220
Specificity	PHX2B Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NBPhox;Neuroblastoma Phox;paired like homeobox 2b;Paired-like homeobox 2B;PMX2B;PHX2B;PHOX2B PMX2B;Paired mesoderm homeobox protein 2B (Neuroblastoma Phox) (NBPhox) (PHOX2B homeodomain protein) (Paired-like homeobox 2B)
Observed Band	34kD
Calculated Molecular Weight	34kD
Cell Pathway	Nucleus .
Tissue Specificity	Expressed in neuroblastoma, brain and adrenal gland.
Function	disease:Defects in PHOX2B are a cause of congenital central hypoventilation syndrome (CCHS) [MIM:209880]; also known as congenital failure of autonomic control or Ondine curse. Most mutations consist of 5-10 alanine expansions in the poly-Ala region from amino acids 241-260. CCHS is a rare disorder characterized by abnormal control of respiration in the absence of neuromuscular or lung disease, or an identifiable brain stem lesion. A deficiency in autonomic control of respiration results in inadequate or negligible ventilatory and arousal responses to hypercapnia and hypoxemia. CCHS is frequently complicated with neurocristopathies such as Hirschsprung disease that occurs in about 16% of CCHS cases.,disease:Defects in PHOX2B predispose to



hereditary neuroblastoma (NB) [MIM:256700]. NB is a tumor of the sympathetic nervous system that account for about 10% of all cancers in childhood. Ger

Background

paired like homeobox 2b(PHOX2B) Homo sapiens The DNA-associated protein encoded by this gene is a member of the paired family of homeobox proteins localized to the nucleus. The protein functions as a transcription factor involved in the development of several major noradrenergic neuron populations and the determination of neurotransmitter phenotype. The gene product is linked to enhancement of second messenger-mediated activation of the dopamine beta-hydroxylase, c-fos promoters and several enhancers, including cyclic amp-response element and serum-response element. Expansion of a 20 amino acid polyalanine tract in this protein by 5-13 aa has been associated with congenital central hypoventilation syndrome. [provided by RefSeq, Jul 2016],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4°over night