



PHC3 Polyclonal Antibody

Catalog No	YP-Ab-05946
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	PHC3 EDR3 PH3
Immunogen	Synthesized peptide derived from human protein . at AA range: 720-800
Specificity	PHC3 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PHC3;PHC3 EDR3 PH3;Polyhomeotic-like protein 3 (Early development regulatory protein 3) (Homolog of polyhomeotic 3) (hPH3)
Observed Band	108kD
Calculated Molecular Weight	108kD
Cell Pathway	Nucleus .
Tissue Specificity	Adipose tissue,Epithelium,Testis,
Function	function:Component of the Polycomb group (PcG) multiprotein PRC1 complex, a complex required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 FCS-type zinc finger.,similarity:Contains 1 SAM (sterile alpha motif) domain.,subunit:Component of chromatin-associated class II polycomb repressive complex 1 (PRC1/hPRC-H) at least composed of PCGF2/RNF110, BMI1/PCGF4, CBX2/M33, CBX4/PC2, CBX8/PC3, PHC1, PHC2, PHC3, SCMH1, RING1 and RNF2/RING2.,

**Background**

function:Component of the Polycomb group (PcG) multiprotein PRC1 complex, a complex required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 FCS-type zinc finger.,similarity:Contains 1 SAM (sterile alpha motif) domain.,subunit:Component of chromatin-associated class II polycomb repressive complex 1 (PRC1/hPRC-H) at least composed of PCGF2/RNF110, BMI1/PCGF4, CBX2/M33, CBX4/PC2, CBX8/PC3, PHC1, PHC2, PHC3, SCMH1, RING1 and RNF2/RING2.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images