



KCNK5 Polyclonal Antibody

Catalog No	YP-Ab-05948
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	KCNK5 TASK2
Immunogen	Synthesized peptide derived from human protein . at AA range: 230-310
Specificity	KCNK5 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Acid-sensitive potassium channel protein TASK-2;K2p5.1;TASK 2;TASK2;TWIK-related acid-sensitive K(+) channel 2;KCNK5;KCNK5 TASK2;Potassium channel subfamily K member 5 (Acid-sensitive potassium channel protein TASK-2) (TWIK-related acid-sensitive K(+) channel 2)
Observed Band	72 kDa
Calculated Molecular Weight	499 aa, 55 kDa
Cell Pathway	Membrane ; Multi-pass membrane protein .
Tissue Specificity	Abundant expression in kidney, also detected in liver, placenta and small intestine. In the kidney, expression is restricted to the distal tubules and collecting ducts (PubMed:9812978). Not expressed in proximal tubules or glomeruli (PubMed:9812978).
Function	function:pH-dependent, voltage insensitive, outwardly rectifying potassium channel. Outward rectification is lost at high external K(+) concentrations., miscellaneous:Inhibited by quinine, quinidine and external acidification., similarity:Belongs to the two pore domain potassium channel (TC 1.A.1.8) family.,subunit:Homodimer; disulfide-linked.,tissue specificity:Abundant expression in kidney, also detected in liver, placenta and small intestine. In the kidney, expression is restricted to the distal tubules and collecting ducts. Not



expressed in proximal tubules or glomeruli.,

Background

This gene encodes one of the members of the superfamily of potassium channel proteins containing two pore-forming P domains. The message for this gene is mainly expressed in the cortical distal tubules and collecting ducts of the kidney. The protein is highly sensitive to external pH and this, in combination with its expression pattern, suggests it may play an important role in renal potassium transport. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images