



ADCY3 Polyclonal Antibody

Catalog No	YP-Ab-06489
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	ADCY3 KIAA0511
Immunogen	Synthesized peptide derived from human protein . at AA range: 950-1030
Specificity	ADCY3 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AC3;AC III;ACIII;AC-III;ADCY 3;ADCY3;ADCY3 KIAA0511;Adenylate cyclase type 3 (EC 4.6.1.1) (ATP pyrophosphate-lyase 3) (Adenylate cyclase type III) (AC-III) (Adenylate cyclase;olfactive type) (Adenylyl cyclase 3) (AC3)
Observed Band	130 kDa,82 kDa
Calculated Molecular Weight	129 kDa
Cell Pathway	Cell membrane ; Multi-pass membrane protein . Cytoplasm . Cell projection, cilium . Golgi apparatus . Also detected in the cytoplasm, close to lipid droplets. .
Tissue Specificity	Detected in zona glomerulosa and zona fasciculata in the adrenal gland (at protein level) (PubMed:11549699). Expressed in brain, heart, kidney, liver, lung, pancreas islets, placenta, and skeletal muscle (PubMed:9920776). Detected in testis (PubMed:15705663).
Function	catalytic activity:ATP = 3',5'-cyclic AMP + diphosphate.,cofactor:Binds 2 magnesium ions per subunit.,enzyme regulation:Activated by calcium/calmodulin.,function:Mediates odorant detection (possibly) via modulation of intracellular cAMP concentration.,similarity:Belongs to the adenylyl cyclase class-4/guanylyl cyclase family.,similarity:Contains 2 guanylate cyclase domains.,tissue specificity:Expressed in brain, heart, kidney, liver, lung, pancreas, placenta, and skeletal muscle.,

**Background**

This gene encodes adenylyl cyclase 3 which is a membrane-associated enzyme and catalyzes the formation of the secondary messenger cyclic adenosine monophosphate (cAMP). This protein appears to be widely expressed in various human tissues and may be involved in a number of physiological and pathophysiological metabolic processes. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Feb 2016]

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images